

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Adolescent and Young Adult (AYA) Oncology

Version 1.2012

NCCN.org

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

[NCCN Guidelines Index](#)
[AYA Oncology Table of Contents](#)
[Discussion](#)

* Peter F. Coccia, MD / Chair € ‡ ξ
UNMC Eppley Cancer Center at
The Nebraska Medical Center

Jessica Altman, MD ‡
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

* Smita Bhatia, MD, MPH € θ
City of Hope Comprehensive
Cancer Center

Scott C. Borinstein, MD, PhD € ‡
Vanderbilt-Ingram Cancer Center

Joseph Flynn, DO, MPH ‡
The Ohio State University
Comprehensive Cancer Center -
James Cancer Hospital and
Solove Research Institute

* Suzanne George, MD †
Dana-Farber/Brigham and Women's
Cancer Center | Massachusetts General
Hospital Cancer Center

Robert Goldsby, MD € ‡
UCSF Helen Diller Family
Comprehensive Cancer Center

* Robert Hayashi, MD € ‡ ξ
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

* Mary S. Huang, MD € ‡
Dana-Farber/Brigham and Women's Cancer
Center | Massachusetts General Hospital Cancer
Center

* Rebecca H. Johnson, MD € Δ
Fred Hutchinson Cancer Research/
Seattle Cancer Care Alliance

* Lynda Kwon Beupin, MD € ‡
Roswell Park Cancer Institute

Michael P. Link, MD €
Stanford Institute Cancer

* Kevin C. Oeffinger, MD ρ θ
Memorial Sloan-Kettering Cancer Center

Kathleen M. Orr, MSW, LCSW-C θ £
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

* Alberto S. Pappo, MD € ‡
St. Jude Children's Research Hospital/
University of Tennessee Cancer Institute

* Damon Reed, MD € ‡
H. Lee Moffitt Cancer Center & Research Institute

* Holly L. Spraker, MD, MS € ‡ θ
Huntsman Cancer Institute
at the University of Utah

Deborah A. Thomas, MD
The University of Texas
MD Anderson Cancer Center

* Margaret von Mehren, MD †
Fox Chase Cancer Center

* Daniel S. Wechsler, MD, PhD € ‡
Duke Cancer Institute

* Kimberly F. Whelan, MD € ‡
University of Alabama at Birmingham
Comprehensive Cancer Center

* Brad Zebrack, MSW, MPH, PhD θ £
University of Michigan
Comprehensive Cancer Center

NCCN

Dorothy A. Shead, MS
Hema Sundar, PhD

Specialties Index:
ξ Bone marrow transplantation
Δ Cancer genetics
‡ Hematology/hematology oncology
ρ Internal medicine
† Medical oncology
€ Pediatric oncology
θ Psychiatry and psychology, including
health behavior
£ Supportive Care including Palliative,
Pain management, Pastoral care and
Oncology social work

Continue

[NCCN Guidelines Panel Disclosures](#)



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 1.2012 Table of Contents Adolescent and Young Adult Oncology

[NCCN Guidelines Index](#)
[AYA Oncology Table of Contents](#)
[Discussion](#)

[NCCN Guidelines AYA Panel Members](#)

[Definition of the AYA Population and Purpose of the AYA Oncology Guidelines \(AYAO-1\)](#)

[Screening, Assessment, and Evaluation \(AYAO-2\)](#)

[Comprehensive Assessment \(AYAO-3\)](#)

[Age-Appropriate Care \(AYAO-4\)](#)

[Treatment-Related Issues \(AYAO-5\)](#)

[Fertility/Endocrine Considerations \(AYAO-6\)](#)

[Psychosocial/Behavioral Considerations](#)

- [Individual \(AYAO-7\)](#)
- [Relationships \(AYAO-8\)](#)
- [Socioeconomic Issues \(AYAO-9\)](#)

[Survivorship \(AYAO-10\)](#)

[Disease-Specific Issues Related to Age \(AYAO-A\)](#)

[Screening Recommendations for AYA Survivors \(AYAO-B\)](#)

[Palliative Care and End-of-Life Considerations \(AYAO-C\)](#)

[Online Resources for AYA Patients and Survivors \(AYAO-D\)](#)

Clinical Trials: NCCN believes that the best management for any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN member institutions, [click here: nccn.org/clinical_trials/physician.html](#).

NCCN Categories of Evidence and Consensus: All recommendations are Category 2A unless otherwise specified.

See [NCCN Categories of Evidence and Consensus](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use, or application, and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2012



DEFINITION OF THE ADOLESCENT AND YOUNG ADULT ONCOLOGY POPULATION

The National Cancer Institute (NCI) Surveillance, Epidemiology, and End Results (SEER) database defines the Adolescent and Young Adult (AYA) Oncology patient as the one diagnosed at 15-29 years of age.^a Subsequently, NCI's AYA Oncology Progress Review Group defines AYA as a patient diagnosed at 15-39 years of age.^b In the NCCN Guidelines, AYA will be defined as patients 15-39 years of age at the time of initial cancer diagnosis.

PURPOSE OF THE NCCN GUIDELINES FOR AYA ONCOLOGY

- These guidelines have been developed as supportive care guidelines and not as treatment guidelines. The purpose of the guidelines is to increase awareness of unique issues in AYA oncology, identify issues and recommend interventions unique to the AYA population. In addition, these guidelines will identify resources available to the AYA population, include appropriate tabular materials, and make recommendations per patient management.
- AYA patients diagnosed with cancer should be recognized as distinct age groups that have unique medical and psychosocial needs. The frequency of distribution of cancer types is dramatically different across the age spectrum of the AYA population.^c
- The distinct biology of disease as well as other age-related issues in the AYA population (fertility, long-term side effects, insurance/financial issues, transportation to clinic appointments, child care, psychosocial support, and adherence to therapy) should be considered in the treatment decision-making process.
- The goal of the NCCN Guidelines for AYA Oncology is to identify issues specific to the AYA population; recommend interventions unique to the AYA population; educate physicians regarding the prevalence of cancer in AYA; its long-term consequences, special considerations related to cancer management in AYA patients that aim to improve treatment tolerance, compliance, and clinical outcomes; promote participation in clinical trials.
- Participation in clinical trials should be strongly encouraged in the AYA population.

^aBleyer A, O'Leary M, Barr R, Ries L. Cancer Epidemiology in Older Adolescents and Young Adults 15 to 29 Years of Age, Including SEER Incidence and Survival: 1975-2000. National Cancer Institute, NIH Pub. No. 06-5767 2006.

^bClosing the Gap: Research and Care Imperatives for Adolescents and Young Adults with Cancer Report of the Adolescent and Young Adult Oncology Progress Review Group. 2006. http://planning.cancer.gov/library/AYAO_PRG_Report_2006_FINAL.pdf


^cFor age-specific incidences rates of cancer by age group and sex in the AYA population, [See Table 2 \(MS-23\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



SCREENING, ASSESSMENT, AND EVALUATION



^dSee [Definition of AYA Population \(AYAO-1\)](#).

^eThese centers provide a multidisciplinary approach involving a team of providers with expertise in cancer treatment and management of specific developmental issues such as fertility, education, career development, employment, family planning, pregnancy, sexually transmitted diseases, smoking, and substance abuse.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

COMPREHENSIVE ASSESSMENT

- Provide age-appropriate information related to cancer
[See Online Resources for AYA Patients and Survivors \(AYAO-D\)](#)
- Discuss risks for fertility and fertility preservation prior to the start of therapy
[See Fertility/Endocrine Considerations \(AYAO-6\)](#)
- Psychosocial assessment
 - ▶ [See Psychosocial/Behavioral Considerations](#)
 - ◊ [Individual \(AYAO-7\)](#)
 - ◊ [Relationships \(AYAO-8\)](#)
 - ◊ [Socioeconomic Issues \(AYAO-9\)](#)
 - ▶ [See NCCN Guidelines for Distress Management](#)
- Genetic and familial risk assessment (within 2 months after the start of therapy)
 - ▶ Risk factors for breast cancer
 - ◊ Germline mutations of *BRCA1*, *BRCA2*, *TP53* (Li-Fraumeni syndrome) or *PTEN* (Cowdens syndrome)
[See NCCN Guidelines for Genetics/Familial High-Risk Assessment: Breast and Ovarian Cancer](#)
 - ◊ Chest irradiation
 - ▶ Risk factors for colon cancer
 - ◊ Mutations in *MMR* genes [hereditary nonpolyposis colorectal cancer (HNPCC or Lynch syndrome)] or *APC* genes [familial adenomatous polyposis (FAP)]
[See NCCN Guidelines for Colorectal Cancer Screening](#)
 - ▶ Risk factors for sarcomas
 - ◊ Li-Fraumeni syndrome
 - ◊ Germline mutations in the retinoblastoma (*RB*) gene or succinate dehydrogenase (*SDH*) gene. Testing for germline mutations in the *SDH* subunit genes should be considered for AYA with wild-type gastrointestinal stromal tumors (GIST) (lacking *KIT* or *PDGFRA* mutations)
 - ◊ FAP-associated desmoid tumors (aggressive fibromatosis)
[See NCCN Guidelines for Colorectal Cancer Screening](#)
 - ▶ Risk factors for malignant peripheral nerve sheath tumors
 - ◊ Germline mutations in neurofibromatosis type I (*NF1*)



Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



AGE-APPROPRIATE CARE

The spectrum of cancer types that affect the AYA population is unique and different from cancer types that affect the pediatric and older populations. When homicide, suicide, and unintentional injury are excluded, cancer is the leading cause of death in this age group. The most common cancers by histologic subtype and by sex are listed below:^{g,h}

Age-appropriate care^f →

Cancer type	Ages 15-39
Females	
Breast carcinoma	20.4
Thyroid carcinoma	14.6
Melanoma	9.5
Carcinoma of cervix and uterus	9.1
HL	3.7
Carcinoma of colon and rectum	3.4
Males	
Gonadal germ cell tumors	10.1
Melanoma	5.5
NHL	4.7
Carcinoma of colon and rectum	3.6
Thyroid carcinoma	2.9

^fThe appropriate location of care and the treatment varies with the type of cancer. Consider referral to medical centers with expertise in treating AYA patients with cancer.

^gRates are per 100,000.

^hData from Howlader N, Noone AM, Krapcho M, Neyman N, Aminou R, Waldron W, Altekruse SF, Kosary CL, Ruhl J, Tatalovich Z, Cho H, Mariotto A, Eisner MP, Lewis DR, Chen HS, Feuer EJ, Cronin KA, Edwards BK (eds). SEER Cancer Statistics Review, 1975-2008, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2008, based on November 2010 SEER data submission, posted to the SEER web site, 2011.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



TREATMENT-RELATED ISSUES

Dose schedules



- AYA patients are usually tolerant of intensive therapies compared to older patients
 - ▶ Dose intensity and dose density are associated with improved outcomesⁱ
 - ▶ [See NCCN Guidelines for Myeloid Growth Factors](#) for growth factor support
- Dose reductions are often based upon avoiding severe, irreversible organ damage
 - ▶ Assume that the patient population has a significant long-term survival and that significant end-organ damage may compromise long term function and quality of life
- Monitoring of cumulative dosing for certain medications associated with irreversible organ damage may be essential when certain life time exposure is encountered
 - ▶ Anthracycline-based chemotherapy (cardiac dysfunction)
 - ▶ Epipodophyllotoxins (secondary acute myeloid leukemia [AML])
 - ▶ Cisplatin (hearing impairment)
 - ▶ Ifosfamide (renal dysfunction)
- Maximum cumulative dosing parameters are often established for a patient to reduce the risk of significant irreversible damage

Toxicities



- Reversible toxicities do not necessarily warrant dose reductions. See NCCN Supportive Care Guidelines for the management of treatment-related toxicities including:
 - ▶ [See NCCN Guidelines for Adult Cancer Pain](#)
 - ▶ [See NCCN Guidelines for Antiemesis](#)
 - ▶ [See NCCN Guidelines for Cancer- and Chemo-Related Anemia](#)
 - ▶ [See NCCN Guidelines for Cancer-Related Fatigue](#)
 - ▶ [See NCCN Guidelines for Palliative Care](#)
 - ▶ [See NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#)
- Intensive screening is recommended for the following treatment-related toxicities:
 - ▶ Cardiac toxicity - Regular echocardiograms (EKG) to monitor cardiac toxicity associated with anthracycline-based chemotherapy
 - ▶ Renal toxicity - Regular glomerular filtration rate (GFR) calculations to monitor renal toxicity associated with cisplatin- and ifosfamide-based chemotherapy
 - ▶ Neurotoxicity - Regular audiogram to monitor hearing loss associated with cisplatin- or carboplatin-based chemotherapy


ⁱWomer RB, West DC, Krailo MD, Dickman PS, Pawel B, for the Children's Oncology Group AEWS0031 Committee. Randomized comparison of every-two-week v. every-three-week chemotherapy in Ewing sarcoma family tumors (ESFT). *J Clin Oncol* (Meeting Abstracts) 2008;26:10501.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



FERTILITY/ENDOCRINE CONSIDERATIONS



^jLevine J, Canada A, Stern CJ. Fertility preservation in adolescents and young adults with cancer. *J Clin Oncol*. 2010;28:4831-4841. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20458029>.

^kThe impact of cancer therapy on fertility is related to the age of the patient at the time of treatment and is dependent on the duration, dose intensity, and type of treatment. [See NCCN Guidelines for Breast Cancer](#) for the management of women with breast cancer during pregnancy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PSYCHOSOCIAL/BEHAVIORAL CONSIDERATIONS

ASSESSMENT

EVALUATION

SUPPORTIVE CARE SERVICES/INTERVENTIONS

- Individual →
- Characteristics
 - ▶ Cognitive function
 - ▶ Emotional issues
([See NCCN Guidelines for Distress Management](#))
 - ▶ Evaluate for other psychiatric symptoms, depression, and anxiety
 - ▶ Living status
 - ◊ Alone
 - ◊ Spouse/partner
 - ◊ Parents
 - Behaviors
 - ▶ Adherence to therapy
 - ▶ Involvement/interruption of school/work
 - ▶ Tobacco, alcohol, or substance abuse
 - ▶ Sexual behavior/risks/concerns
 - ▶ Dietary needs
 - ▶ Exercise needs
 - Existential/spiritual issues

Relationships → [See AYA0-8](#)

Socioeconomic issues → [See AYA0-9](#)

- Refer AYA patients with cognitive dysfunction or other psychiatric symptoms (eg, depression, or anxiety) to a mental health provider and community-based resources serving AYA patients
- Offer psychosocial support and counseling to help alleviate distress.
[See NCCN Guidelines for Distress Management](#)
- Adherence to therapy
 - ▶ Provide education and/or guidance about each medication prior to the start of treatment and every time there is a change in treatment.
 - ▶ Review list of medications and their dose, purpose, and adverse effects.
 - ▶ Simplify dosing schedule and change timing and frequency of medication or method of administration, when medically possible, to fit into AYAs lifestyle and normal activities.
 - ▶ Provide access to systematic and standardized symptom management for side effects related to cancer treatment.
[See NCCN Guidelines for Supportive Care](#)
- Provide flexible treatment dates, consultation times, and procedures to enable AYAs to continue with their treatment without interrupting their normal activities (school/work).
- Refer patients with signs, symptoms, and a history of substance abuse or addiction to a risk reduction or substance abuse management program
- Provide health education about sexually transmitted infections, diet, and exercise.
- For all AYA patients, provide counseling regarding the risks of treatment-related infertility and discuss options for fertility preservation prior to the start of therapy. [See Fertility/Endocrine Considerations \(AYA0-6\)](#)
- Refer patients experiencing challenges with their faith or belief in a just or fair world to faith-based resources or activities (eg, church youth groups, mentors)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PSYCHOSOCIAL/BEHAVIORAL CONSIDERATIONS

ASSESSMENT

EVALUATION

SUPPORTIVE CARE SERVICES/INTERVENTIONS

Relationships →

- Family status
 - ▶ Interaction and relationship with parents
 - ▶ Interaction and relationship with spouse/partner
- Peer relationships
- Participation in community and social activities (eg, religious organizations, clubs, athletics/recreation, music, youth groups)
- Communications with health care professionals
 - ▶ Decision-making preferences: parents, clinical, and/or self
 - ▶ Information and communication preferences (eg, parents)

→

- Promote communication between AYA patients and family members
 - ▶ Parents
 - ▶ Spouse/partners
 - ▶ Siblings
- Provide family members and partners with information about psychosocial support and behavioral services
 - ▶ Increase awareness of the possible psychosocial issues associated with cancer diagnosis in AYAs, so that family members and partners may continue to support the patient.
- Family-based intervention models from pediatric studies may have utility for AYAs
 - ▶ Parent support groups
 - ▶ AYA support groups
 - ▶ Social and recreational programs
 - ▶ Psychoeducational programs
- Provide information about peer-support groups to assist AYAs establishing and maintaining relationships with their normal peers as well as with other AYAs with cancer.
 - ▶ [See Online Resources for AYA Patients and Survivors \(AYAO-D\)](#)
 - ▶ Face-to-face meetings
 - ▶ Camp and retreat programs
 - ▶ Online support groups
- Create flexible visiting hours and an environment that will encourage peers to visit AYA patients
- Health care professionals should establish direct communication with individual patients
 - ▶ Reinforce the importance of AYA involvement in decision-making
 - ▶ Provide age-appropriate information about their cancer, treatment options, and potential side effects
 - ▶ Ask for permission to share information with family members

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PSYCHOSOCIAL/BEHAVIORAL CONSIDERATIONS

ASSESSMENT

EVALUATION

SUPPORTIVE CARE SERVICES/INTERVENTIONS

Socioeconomic issues →

- Insurance availability and security
 - ▶ Employer -provided
 - ▶ Parent's insurance
- Assessment of risk for losing insurance
 - ▶ Loss of employment
 - ▶ Age out of parents' insurance
- Risk for financial loss or bankruptcy
- Child care
- Transportation
- Accommodation if traveling to receive treatment
- Desires for complementary and alternative medicine (CAM)

- Link qualified AYA patients to Medicaid, social security, and/or disability insurance
- Educate AYA patients about benefits they may qualify for, such as short- or long-term disability, state disability benefits, and/or food stamps.
- Direct AYA patients to legal resources/advocates for understanding health insurance coverage
- Identify resources for respite care for AYA patients with young children
- Refer to transportation assistance programs (eg, van ride programs; voucher programs)
- Refer to reputable providers of CAM services
- Provide AYAs with a list of recommended and reliable online sources to access information related to their cancer
[See Online Resources for AYA Patients and Survivors \(AYAO-D\)](#)
- Financial assistance for AYA cancer survivors needs to be integrated into survivorship plans.
- AYAs with cancer need long-term follow-up care for monitoring and treatment of late effects long after completion of treatment.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2012


Adolescent and Young Adult Oncology

SURVIVORSHIP

AYA cancer survivor¹ →

- Develop a “Cancer Treatment Summary and Survivorship Care Plan”
- Provide a periodic evaluation focusing on history, physical examination and screening based upon treatment exposures, and risk for treatment-related late effects
- Counsel regarding lifestyle practices and methods to reduce risk (eg, avoiding smoking, increasing level of physical activity)
- Advocate for appropriate health care coverage
- Recommend a dental exam and cleaning every 6 months

SELECTED EXPOSURES



SCREENING RECOMMENDATIONS

[See Screening Recommendations \(AYAO-B\)](#)

[See AYAO-B](#)

¹AYA cancer survivorship occurs after successful completion of therapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



DISEASE-SPECIFIC ISSUES RELATED TO AGE

Acute lymphoblastic leukemia (ALL)

- See NCCN Guidelines for ALL (coming March 2012)

Bone and soft tissue sarcomas

- [See NCCN Guidelines for Bone Cancer](#) and [NCCN Guidelines for Soft Tissue Sarcoma](#)
- Rhabdomyosarcoma
 - ▶ Uncommon outside of the pediatric population; should be referred to an institution with experience in the management of rhabdomyosarcoma

Colon cancer

- Higher incidence of mucinous histology
- More often right-sided
- Higher incidence of signet ring cells and microsatellite instability (MSI)
- More advanced stage at diagnosis
- Lower incidence of *KRAS* mutations
- Decreased incidence of chromosomal instability
- Consider mismatch repair gene deficiency in these patients
- Increased risk for additional malignancies

Melanoma

- Melanocytic tumors of uncertain malignant potential (MELTUMP) are more frequently seen in younger patients and when suspected, referral to a pathologist with expertise in atypical melanocytic lesions is recommended.
- Principles of pathology for younger patients with consideration to additional testing comparative genomic hybridization (CGH) or fluorescent in situ hybridization (FISH) may be useful to detect the presence of selected gene mutations for histologically equivocal lesions. [See NCCN Guidelines for Melanoma](#).
- Sentinel lymph node biopsy
 - ▶ Higher yield in AYA population
- Surgical margins have not been established for patients <18 years old as they were not included in the trials.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



SCREENING RECOMMENDATIONS FOR AYA SURVIVORS

The following screening recommendations are adapted from the treatment exposure-based COG Long-Term Follow-Up Guidelines for Survivors of Childhood, Adolescent, and Young Adult Cancers.

- The recommendations represent only key aspects; for more detail, refer to the website survivorshipguidelines.org.
- The COG Guidelines are based upon exposures used in the treatment for pediatric cancer. As such, the recommendations are applicable to many survivors of cancers that span across adolescence and young adulthood, such as acute leukemias, Hodgkin's and non-Hodgkin's lymphomas, medulloblastomas, and sarcomas. In addition, since the treatment exposures for some young adult cancers, such as male germ cell tumors, are similar to pediatric cancer treatments (eg, cisplatin, bleomycin, abdominal irradiation), the recommendations may be applicable. In contrast, the COG recommendations are generally not applicable to survivors of typical adult carcinomas occurring during young adulthood, such as breast, colorectal, and ovarian cancers.
- The risk for many late effects may be influenced by family history, lifestyle behaviors, and comorbid health conditions. The following recommendations are based upon the treatment exposure; timing and intensity of screening may be adapted based upon additional risk factors.
- Most survivors will have multiple treatment exposures, and therefore may have multiple screening needs.

Neuroendocrine axis screening (selected outcomes)

- Growth hormone deficiency
 - High-risk population: radiation dose to hypothalamic-pituitary-adrenal (HPA) axis >18 Gy

- Screening recommendation: height, weight, and body mass index every 6 months until growth is completed then yearly. Note: most AYA patients will have attained (or nearly attained) final height; the significance and management of growth hormone status among survivors who attained their final height is controversial
- Consider endocrine consultation for height below the third percentile on the growth curve, drop of less than second percentile rankings on the growth chart
- Central hypothyroidism
 - High-risk population: total radiation dose to HPA axis >40 Gy
 - Screening recommendation: thyroid-stimulating hormone (TSH) and free T4, yearly
- Gonadotropin deficiency
 - High risk population: total radiation dose to HPA axis >40 Gy
 - Screening recommendation: FSH, luteinizing hormone (LH), and testosterone (males) and FSH, LH, and estradiol (females) as clinically indicated; semen analysis (males) as requested by patient or for evaluation of fertility
- Central adrenal insufficiency
 - High-risk population: total radiation dose to HPA axis >40 Gy
 - Screening recommendation: 8:00 AM. serum cortisol, yearly for at least 15 years after treatment and as clinically indicated

Neuropsychological evaluation

- Severe neurocognitive deficits are uncommon in survivors of AYA cancer, including CNS tumors. However, subtle deficits in executive function, sustained attention, memory, and processing speed may occur with higher dose cranial radiation (>18 Gy)
- Screening recommendation: In patients with evidence of impaired educational or vocational progress, formal neuropsychological evaluation is recommended

[Continued on next page](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



SCREENING RECOMMENDATIONS FOR AYA SURVIVORS

Breast cancer screening (females)

- High-risk population: chest radiation >20 Gy prior to the age of 30 years
- Screening recommendation: breast magnetic resonance imaging (MRI) and mammogram yearly, starting at age 25 or 8 years after radiation, whichever occurs last

Cardiovascular risk assessment and screening

- High risk populations: TBI, mediastinal/chest radiation >20 Gy
- Screening recommendation: measure blood pressure and body mass index yearly; fasting glucose, lipid profile every two years
- Screening for ischemic coronary artery disease remains controversial; consider cardiology consultation (5 to 10 years after radiation) in patients who received >40 Gy chest radiation

Screening for cardiomyopathy/asymptomatic heart failure

- High-risk population: cumulative anthracycline dose >300 mg/m²; chest radiation >30 Gy; combination of anthracycline and chest radiation
- Screening recommendation: EKG (or MUGA scan) every 1-2 years (Note: Frequency of testing is dependent on both age at time of exposure and dose of exposure. The frequency of testing has not been established for breast cancer survivors treated with lower cumulative doses of anthracyclines.)

Screening for valvular heart disease

- High risk population: chest radiation >30 Gy
- Screening recommendation: EKG every 1-2 years

Pulmonary screening

- High-risk population: chest radiation >15 Gy (or radiation to large volume of lung) TBI (>6 Gy in single fraction or >12 Gy fractionated), bleomycin >400 U/m², combination of chest radiation and bleomycin, and selected alkylating agents (busulfan >500 mg, carmustine >600 mg/m²)
- Screening recommendation: chest x-ray and pulmonary function tests (including diffusion lung capacity for carbon monoxide [DLCO] and spirometry) as a post-therapy baseline and then as clinically indicated in patients with abnormal results or progressive pulmonary dysfunction

Thyroid screening

- Thyroid disorders: hypothyroidism (very common), thyroid cancer (common), and hyperthyroidism (uncommon)
- High-risk population: radiation field includes the thyroid gland (see neuroendocrine axis screening for high-dose cranial radiation)
- Screening recommendation: TSH and thyroid/neck exam, yearly

Colorectal cancer screening

- High-risk population: abdominal or pelvic radiation >30 Gy
- Screening recommendation: colonoscopy starting at age 35 or 10 years after radiation, whichever occurs last

Screening for kidney and/or bladder disease

- Renal insufficiency and secondary renal/renovascular hypertension
 - ▶ High-risk population: radiation >10 Gy, combination of radiation with nephrotoxic agents (eg, cisplatin, ifosfamide, aminoglycosides, amphotericin, immunosuppressants)
 - ▶ Screening recommendation: post-therapy baseline blood urea nitrogen (BUN), creatinine, Na, K, Cl, CO₂, Ca, Mg, PO₄; repeat as clinically indicated; measure blood pressure yearly, and urinalysis yearly

[Continued on next page](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



SCREENING RECOMMENDATIONS FOR AYA SURVIVORS

- Hemorrhagic cystitis/bladder fibrosis
 - ▶ High risk population: cyclophosphamide >3 gm/m², pelvic radiation >30 Gy
 - ▶ Screening recommendation: urinalysis, yearly
- Bladder cancer
 - ▶ High risk population: cyclophosphamide combined with pelvic radiation
 - ▶ Screening recommendation: urinalysis, yearly

Assessment for gonadal function

- Males
 - ▶ Infertility
 - ◊ High-risk population: moderate- to high-dose alkylating agent chemotherapy (eg, MOPP >3 cycles, Busulfan >600 mg/m², cyclophosphamide cumulative dose >7.5 gm/m² or as conditioning for hematopoietic cell transplant, ifosfamide cumulative dose >60 gm/m²), TBI, testicular irradiation >2 Gy, and any alkylator combined with testicular irradiation or TBI
 - ◊ Screening recommendation: semen analysis as requested by patient or for evaluation of infertility; periodic evaluation over time is recommended as resumption of spermatogenesis can occur up to 10 years post therapy
 - ▶ Leydig cell dysfunction
 - ◊ High-risk population: testicular irradiation >20 Gy
 - ◊ Screening recommendation: testosterone as clinically indicated in patients with clinical signs and symptoms of testosterone deficiency

- Females
 - ▶ Infertility (acute ovarian failure or premature menopause)
 - ◊ High-risk population: moderate- to high-dose alkylating agent chemotherapy (eg, MOPP >3 cycles, Busulfan >600 mg/m², cyclophosphamide cumulative dose >7.5 gm/m² or as conditioning for hematopoietic cell transplant, ifosfamide cumulative dose >60 gm/m²), TBI, and abdominal and/or pelvic radiation
 - ◊ Screening recommendation: FSH, LH, estradiol as indicated in patients with irregular menses, primary or secondary amenorrhea, and/or clinical signs and symptoms of estrogen deficiency

Screening for t-AML or myelodysplasia

- High risk populations: epipodophyllotoxins, alkylating agents, cisplatin, and/or anthracyclines
- Screening recommendation: Complete blood count (CBC)/differential yearly, up to 10 years after exposure

Audiological evaluation

- High-risk population: cisplatin >360 mg/m², radiation involving the ear >30 Gy, and combination of cisplatin and cranial/ear radiation
- Screening recommendation: audiology testing as a post-therapy baseline and then as clinically indicated for signs and symptoms of hearing loss

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

**PALLIATIVE CARE/END-OF-LIFE CONSIDERATIONS**

Palliative care was commonly called “end-of-life care” but it is now accepted as a treatment to control symptoms, and reduce emotional and physical suffering at any stage of a life-threatening disease. Referral to palliative care is still appropriate when patients are being treated with curative intent. The World Health Organization (WHO) definition of palliative care includes palliative care being initiated at diagnosis.¹ A palliative care team should involve a multidisciplinary team with expertise in understanding the psychosocial, emotional, and developmental issues that are unique to the AYA population.² [See NCCN Guidelines for Palliative Care](#)

Four main ideas of palliative care needs for AYA population:¹

- Psychosocial needs^{3,4}

[See Psychosocial/Behavioral Considerations \(AYAO-8\)](#)

- ▶ Psychosocial needs of the patient
 - ◊ Needs depends on maturity, and level of independence (loss of new-found independence)
 - ◊ Peer support - Facilitate peer relationships and interaction with patient’s normal peers as well as other AYAs with cancer
 - ◊ Provide physical space in a clinic, hospital, etc for social interactions, web-based peer support, and social networking
 - ◊ Provide age-appropriate professional psychosocial support services
- ▶ Psychosocial needs of family and friends
 - ◊ Grief from loss of loved ones begins even before death. Provide family members and friends with information about palliative care services
 - ◊ Provide regular counseling and psychosocial support
- ▶ Psychosocial needs of the professional caregivers
 - ◊ There is a high rate of “burnout” among palliative caregivers
 - ◊ Provide support for debriefing and to maintain a balance between home and work

- Introducing palliative care

- ▶ Lack of communication about illness trajectory is a barrier for transition to palliative care
- ▶ Introduction of palliative care for symptom management and psychosocial support should occur before the patient is considered “palliative” in order to provide the best possible care for the patient
- ▶ Efforts should be made to normalize palliative care involvement without providing a negative connotation of terminal care.

- Resources required

[See Psychosocial /Behavioral Considerations \(AYAO-9\)](#)

- ▶ Insurance availability and security- Provide information about Medicaid, social security, and disability insurance
- ▶ Provide flexibility in the health care system for patients to maintain “normalcy”
- ▶ Evidence-based guidelines in AYA palliative care are limited and need to be developed

- Advocacy

- ▶ AYA specific advocacy groups need to be developed at the state/national/international level to increase awareness.
- ▶ It is important to create an AYA team that includes palliative care in order to improve early referrals, research, and patient-centered care.²

[Continued on next page](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PALLIATIVE CARE and END-OF-LIFE CONSIDERATIONS

End-of-life considerations

- **AYAs understand that death is permanent and irreversible. It is imperative for health care professionals not to assume that AYA patients may be less inclined to discuss death and other end-of-life issues.²**
- **Adolescents indicate a preference for dying at home, yet 80% have died in a hospital.^{5,6}**
- **Palliative care physicians should facilitate discussion about end-of-life care issues such as nutrition/hydration, sedation treatment cessation and place of death.²**
- **An advanced care planning document may be appropriate and helpful for terminally ill AYA patients with metastatic cancer⁷**

¹Pritchard S, Cuvelier G, Harlos M, Barr R. Palliative care in adolescents and young adults with cancer. *Cancer*. 2011;117:2323-2328.

²Wein S, Pery S, Zer A. Role of palliative care in adolescent and young adult oncology. *J Clin Oncol*. 2010;28:4819-4824.

³D'Agostino NM, Penney A, Zebrack B. Providing developmentally appropriate psychosocial care to adolescent and young adult cancer survivors. *Cancer*. 2011;117:2329-2334.

⁴Zebrack BJ. Psychological, social, and behavioral issues for young adults with cancer. *Cancer*. 2011;117:2289-2294.

⁵Bell CJ, Skiles J, Pradhan K, Champion VL. End-of-life experiences in adolescents dying with cancer. *Support Care Cancer*. 2010;18:827-835.

⁶Webb NM, Tucker D. Young adults' opinions about hospice and home death. *J Palliat Med*. 2009;12:337-342.

⁷Wiener L, Ballard E, Brennan T, et al. How I wish to be remembered: the use of an advance care planning document in adolescent and young adult populations. *J Palliat Med*. 2008;11:1309-1313.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

NCCN Guidelines Version 1.2012
Adolescent and Young Adult Oncology**ONLINE RESOURCES for AYA PATIENTS and SURVIVORS**

Website	URL
15-40 Connection	http://www.15-40.org/
AYA Cancer Patients and Survivors: Teenage, Adolescent and Young Adult	http://www.facebook.com/ayacancer
Cancer care Support Groups	http://www.cancercare.org/support_groups
First Descents (Free outdoor adventure programs for young adult cancer fighters and survivors ages 18 to 39)	http://firstdescents.org/
Focus under Forty	http://university.asco.org/focusunder40
Gilda's Club	http://www.gildasclub.org/
I'm Too Young For This! Cancer Foundation	http://stupidcancer.com/about/index.shtml
Imerman Angels (One-on-One Cancer support-Connecting Cancer Fighters, Survivors and Caregivers)	http://www.imermanangels.org/
LIVESTRONG Young Adult Alliance	http://www.livestrong.org/LIVESTRONG-Young-Adult-Alliance
Mass Kickers Foundation	http://masskickers.org/
National Coalition for Cancer Survivorship	http://www.canceradvocacy.org/
Planet Cancer	http://www.planetcancer.org/
Surviving and Moving Forward: The SAMFund for Young Adult Survivors of Cancer	http://www.thesamfund.org/
Teens Living With Cancer	http://www.teenslivingwithcancer.org/
The Jack & Jill Late Stage Cancer Foundation	http://www.jajf.org/
The Oncofertility Consortium	http://oncofertility.northwestern.edu/
Ulman Cancer Fund for Young Adults	http://www.ulmanfund.org/
We Believe Foundation	http://www.webelievefoundation.com
Young Survival Coalition (for young women diagnosed with breast cancer)	http://www.youngsurvival.org/

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

Discussion

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Overview

Over the past 20 years, advances in cancer treatment have significantly improved survival rates for young children and older adults, but there has been no significant improvement in adolescent and young adult (AYA) patients with cancer.¹ One of the main reasons for the lack of improvement in outcomes is that AYA patients have a low rate of participation in clinical trials.¹⁻³ In the United States, approximately 10% of patients 15 to 19 years of age and 1%-2% of patients 20 to 39 years of age are enrolled in clinical trials.⁴ In addition to the low rate of participation in clinical trials, several other factors also contribute to the poor outcome in AYA patients with cancer, such as: differences in disease biology, lack of consistency in treatment approaches, poor compliance with or intolerance of therapy, lack of health insurance, delays in diagnosis, and physician's lack of familiarity with cancer in the AYA population.⁵

The biology, epidemiology, and clinical outcomes affecting AYAs with cancer are usually different than those of older individuals with cancer.⁶ In addition, the genetic, physiological, and pharmacological changes associated with the AYA population may impact AYAs' ability to tolerate cancer therapy and response to treatment. Unlike comprehensive geriatric assessment, which is helpful to physicians in developing a coordinated treatment plan and understanding the functional needs of older patients, no similar assessment has been reported for AYA patients. There are less evidence-based data to guide the treatment of these patients. AYA patients diagnosed with cancer should be recognized as a distinct age group that has unique medical and psychosocial needs.⁷ The distinct biology of disease as well as age-related issues in the AYA population should be considered in the treatment decision-making process.

The AYA patient is generally defined as an individual 15 to 39 years of age at the time of initial cancer diagnosis, although variations may be appropriate depending on individual malignancies.⁸ Nearly 70,000 patients in this age group are diagnosed with cancer each year in the United States, over seven times more patients than those diagnosed who are less than 15 years of age (Table 1). The spectrum of cancer types that affect the AYA population is unique and different from those that affect the pediatric and older population. Cancer is the leading cause of death among the AYA population, excluding homicide, suicide, or unintentional injury.^{5,9} Lymphomas, melanoma, testicular cancer, female genital tract malignancies, thyroid cancer, bone and soft tissue sarcomas, leukemias, central nervous system (CNS) cancers, breast cancer, and non-gonadal germ cell tumors account for 95% of the cancers in this age group.¹ The frequency and incidence of distribution of cancer types is also dramatically different across the age spectrum of



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

the AYA population. See Table 2 for age-specific SEER incidences of cancer by age group and sex in the AYA population.¹⁰

Quality care for AYA patients with cancer is tied to timely detection, efficient processes for diagnosis, initiation of treatment, promotion of adherence, and access to a multidisciplinary team of health care professionals who are well-versed in the specific issues relevant to this patient population. These issues include fertility, long-term side effects, psychosocial and socioeconomic issues, transportation to clinic appointments, child care, treatment adherence, and the unique biology of disease.¹¹

The goal of the NCCN Guidelines for AYA Oncology is to identify issues specific to the AYA population and recommend interventions unique to this patient population; educate physicians regarding the prevalence of cancer in AYA and its long-term consequences; identify special considerations related to the management of cancer in AYA patients with the aim of improving treatment tolerance, compliance, and clinical outcomes; and promote participation in clinical trials.

Risk factors

With rare exceptions, cancer appears to arise sporadically in most AYAs with a negative family history of cancer. There are no established risk factors for the majority of cancer diagnoses before the age of 30.⁹ Toxic and environmental exposures that cause cancer in AYAs include chemotherapy and/or radiation therapy (RT) leading to second malignancies in patients treated for cancer during childhood or young adulthood; predisposition to clear cell adenocarcinoma of the vagina or cervix in patients with maternal exposure to diethylstilbestrol; and melanomas induced by ultraviolet light. Infections that predispose AYAs to cancer include cervical carcinoma following exposure to human papillomavirus (HPV), Hodgkin's lymphoma (HL) and Burkitt's

lymphoma following Epstein-Barr virus (EPV) infection, and Kaposi sarcoma and non-Hodgkin's lymphoma (NHL) in patients with human immunodeficiency virus (HIV).⁹

Familial cancer syndromes, associated with germline mutations in a variety of genes, affect only a small minority of AYA patients with cancer. However, these syndromes greatly increase the risk for cancer during adolescence and young adulthood for affected patients.

Breast cancer

Young women with germline mutations of *BRCA1*, *BRCA2*, *TP53* (Li-Fraumeni syndrome), or *PTEN* (Cowden's syndrome), or those who have received mantle field irradiation for HL are at an increased risk of developing breast cancer during young adulthood.^{1,11} Screening for breast cancer may be warranted in AYA patients with inherited or familial risk factors. See the NCCN Guidelines for Genetics/Familial High-Risk Assessment: Breast and Ovarian Cancer Screening.

Colon cancer

In young adults, hereditary polyposis and nonpolyposis syndromes, inflammatory bowel disease, and radiation exposure are predisposing factors for developing colorectal cancer.¹ Hereditary nonpolyposis colorectal cancer (HNPCC or Lynch syndrome) is an autosomal dominant syndrome caused by mutations in one of the four *MMR* genes (*MSH2*, *MLH1*, *MSH6*, or *PMS2*), and is associated with colon cancer developing in the AYA population.¹² Familial adenomatous polyposis (FAP) is an autosomal dominant disease caused by germline mutations in the *APC* gene. This syndrome is associated with thousands of colonic polyps and with the development of colon cancer in most affected patients by age 40. Desmoid tumors are considered to be the most common extracolonic manifestations of FAP, and may be the presenting manifestation of FAP in AYA patients.¹² Screening for



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

colorectal cancers may be warranted in AYA patients with inherited or familial risk factors. See the NCCN Guidelines for Colorectal Cancer Screening.

Sarcomas

AYA patients with Li-Fraumeni syndrome or germline mutations in the retinoblastoma (*RB*) gene are at a higher risk of developing osteosarcoma. AYA individuals with germline mutations in the *RB* gene have often been treated for RB during early childhood.¹³ AYAs with a family history of Li-Fraumeni syndrome have a higher risk of developing not only sarcomas but a wide variety of malignancies, including leukemia, brain tumors, breast cancer, and adrenocortical carcinoma before 40 years of age.¹⁴ Patients with succinate dehydrogenase (*SDH*) gene mutations are at risk for paraganglioma and pheochromocytoma, gastrointestinal stromal tumors (GISTs), renal clear cell carcinoma, and papillary thyroid carcinoma during adolescence and young adulthood. Testing for germline mutations in the *SDH* subunit should be considered for AYA patients with wild-type GISTs lacking *KIT* or *PDGFRA* mutations.^{15, 16} Patients with germline mutations in neurofibromatosis type I (*NF1*) carry a 10% lifetime risk for malignant peripheral nerve sheath tumors, as well as an increased risk for other malignancies including GISTs and early breast cancer in females.¹⁷

Screening

Cancer screening in some circumstances, particularly in cervical, breast, and colorectal cancers, can significantly reduce mortality if directed at the appropriate age group and if the results are interpreted and followed up appropriately.¹⁸ However, there are no age-specific screening tests that have been developed that would increase early detection in AYAs with cancer, and in some instances screening tests have been associated with false-positive results leading to false

diagnosis and unnecessary treatments.¹⁹ Therefore, it is necessary to identify simple and accurate tests as well as potential risks and benefits before implementing patients in the AYA population. AYAs with cancer should be made aware of the importance of early diagnosis and self-examination of the skin, breast (for females), and testicles (for males) as recommended by the American Cancer Society (ACS). They should also be educated regarding the benefits of early detection and treatment.⁵

Diagnosis

The onset of new symptoms in AYAs may not immediately trigger evaluation for malignancy, due to the relatively low incidence of cancer in this age group and the resulting low index of suspicion on the part of patients and primary care providers. AYAs are at a higher risk of delayed cancer diagnosis, which may result in a more advanced stage of cancer that requires more therapy and is associated with a worse prognosis.⁵ Some studies have reported that adolescents experience longer lag times (interval between symptom onset and diagnosis) than children.²⁰⁻²² Lack of health insurance, inexperienced physicians, and workup that is inappropriate for the patient's age are some causes of delayed diagnosis in AYAs with cancer. In a retrospective analysis of 503 patients aged 15 to 29 years with previously untreated cancer, the advanced stage of cancer at diagnosis and lack of health insurance were significantly associated with longer lagtimes.²³ Those with public or no health insurance had longer lag times than those with private health insurance in most of the cancers evaluated. Patients with leukemia and NHL had shorter lag times (2 to 5 weeks) than those with sarcomas and thyroid cancer (20 to 24 weeks), irrespective of the insurance type. In addition to health insurance, education and employment status are also likely to influence lag time, although these factors were not evaluated in this study.

Special considerations in the management of AYA patients with cancer

All AYA patients should undergo comprehensive assessment following the diagnosis of cancer, which should include psychosocial assessment, discussion of risks of infertility associated with treatment and options for fertility preservation, and genetic and familial risk assessment (within 2 months after the start of therapy).

Age-appropriate care: Pediatric vs. adult cancer centers

AYA patients with cancer can be treated either at pediatric or at adult cancer centers.²⁴ Retrospective analyses have shown that AYA patients with certain pediatric-type cancers, such as acute lymphoblastic leukemia (ALL),²⁵⁻²⁹ rhabdomyosarcoma,³⁰ and Ewing's sarcoma,³¹ have superior outcomes when treated with pediatric protocols. Alternatively, there is a lack of compelling evidence that pediatric protocols improve outcomes in AYA patients with acute myeloid leukemia (AML), HL, and NHL.³²⁻³⁴

As mentioned earlier, the low rate of participation in clinical trials is one of the main reasons for the lack of improvement in outcomes in AYA patients with cancer.¹⁻³ Care should be provided at medical centers with broad access to clinical trials (standard-of-care registry trials and trials evaluating novel therapies).²⁴ Pediatric cancer centers enroll more adolescents into clinical trials (35% vs. 12% at non-pediatric cancer centers), and AYA patients treated at pediatric cancer centers have a higher rate of clinical trial enrollment (26%) compared to those treated at adult cancer centers (4%).³⁵⁻³⁷ More recently, Parsons et al. reported that AYA patients who are treated by non-pediatric oncologists are less likely to be enrolled in clinical trials.³⁸ Nevertheless, a substantial number of AYA patients with pediatric malignancies are not being treated at pediatric cancer centers.^{39,40}

The treatment and appropriate location of care vary with the type of cancer as well as with the availability of family, community, and institutional supports.^{5,41} Most importantly, AYA patients should be evaluated at medical centers with extensive experience in treating cancer in this patient population and at centers that have access to supportive care services (psychosocial/educational support and fertility preservation) specific to AYA population as well as medical subspecialty services appropriate to the cancer diagnosis, such as orthopedic surgeons with experience in limb-sparing surgery for patients with extremity sarcomas.²⁴ Centers should adopt the appropriate evidence-based approach, which includes adult centers implementing treatment based on pediatric protocols that have demonstrated superior outcomes in AYA patients and pediatric centers adopting adult regimens that have demonstrated benefit in this patient population.

AYA patients should be managed by a multidisciplinary team of providers with expertise in cancer treatment and management of specific developmental issues such as fertility, education, career development, employment, family planning, pregnancy, sexually transmitted diseases, and tobacco, alcohol, and substance abuse. Given the rarity of several tumor types diagnosed in this population, all AYA patients should be offered and encouraged to participate in tumor banking studies and multicenter clinical trials, when available.

Treatment options

AYA patients can usually tolerate more intensive therapies than older adults, since they have fewer comorbid conditions that limit the intensity of treatment in older adults. Dose-intensive and dose-dense treatment is associated with improved outcomes. Every AYA patient with cancer should be treated with aggressive therapy if there are no contraindications.²⁹



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

Treatment-related issues in AYA patients with cancer may differ from those of pediatric or older adult patients due to the distinct biology of the disease.⁶ Physical and physiological changes, such as changes in body composition, size and maturity of organs, and hormones associated with the normal pubertal process, may directly affect the drug disposition, drug efficacy, and toxicity of chemotherapy in AYAs.⁴² As mentioned above, AYA patients have fewer comorbid conditions compared to older cancer patients, and thus are usually able to tolerate intense chemotherapy and surgery with less morbidity. Appropriate management of symptoms and side effects to reduce the severity and toxicity of treatment should be an integral part of the management of AYAs with cancer.⁴³

Surgery, RT, chemotherapy, and hematopoietic stem cell transplantation (HSCT) are the main treatment options for patients who are able to tolerate curative treatment. All of these options are associated with both acute and late side effects.^{5, 44}

Surgery

Surgery is more feasible in AYAs with cancer since they have less comorbidities than older patients and anesthesia is easier to administer than in children.⁵ At the same time adolescent patients, whose bodies are still developing, may be more affected by some surgical procedures than older patients who are already at or near their full body size. The extent of surgery is dependent on the type and location of cancer. In some cases, extensive surgery requiring removing part or all of an organ or limb may be necessary. With the recent advances in surgical techniques and chemotherapy, limb-sparing surgery is now feasible for the majority of patients with extremity sarcoma and osteosarcoma.^{3,43}

Radiation therapy

RT has been associated with an increased risk for late mortality; development of second malignancies; pulmonary, cardiac, and thyroid dysfunction; and chronic health conditions and growth abnormalities.⁴⁵ AYAs with cancer receiving RT to testes or ovaries are at risk of developing infertility later in life. Women who receive chest radiation for HL between 10 and 30 years of age are at increased risk of developing breast cancer.⁴⁶ Cranial RT is associated with short stature, cognitive processing difficulties, and poor physical function, which contribute to lower rates of employment, independent living, and marriage among AYA cancer survivors.⁴⁷ See also the sections on “Impact of treatment on fertility and fertility preservation” and “Late effects in AYA cancer survivors.” Adolescents are more vulnerable to radiation-induced spinal cord dysfunction, presumably because of elongation of the cord during the growth spurt.⁴⁸

Chemotherapy

Alkylating agent-based chemotherapy is associated with a higher risk of infertility in both male and female patients.⁴⁶ See the section on “Impact of treatment on fertility and fertility preservation.” Anthracycline-based chemotherapy is associated with cardiac dysfunction, whereas neurotoxic chemotherapies such as methotrexate and cytarabine can result in CNS dysfunction.⁴³ Higher cumulative doses of cisplatin, ifosfamide, or epipodophyllotoxin are associated with hearing loss, renal dysfunction, and secondary AML, respectively.⁴⁹⁻⁵² See also the section on “Late effects in AYA cancer survivors.”

Pain, fatigue, nausea, vomiting, mucositis, hair loss, infection, and myelosuppression are some of the acute side effects of chemotherapy. Reversible toxicities (as mentioned above) do not necessarily warrant dose reductions. See the NCCN Guidelines for Supportive Care for the management of treatment-related toxicities. Every attempt should be



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

made to maintain dose intensity unless it is contraindicated. Dose reductions are often based upon avoiding severe, irreversible organ damage. Significant end-organ damage may compromise long-term function and quality of life in AYA patients. Maximum cumulative dosing parameters are often established for a patient to reduce the risk of significant irreversible damage. Monitoring of cumulative dosing and intensive screening is essential for patients receiving chemotherapy regimens associated with irreversible organ damage.

Anticipatory nausea and vomiting (ANV), also known as conditioned, learned, or psychological nausea and vomiting, is reported to occur before chemotherapy in approximately 20% of patients at any one chemotherapy cycle and in 25%-30% of patients by their fourth chemotherapy cycle.⁵³ Younger patients (less than 50 years of age) may be more susceptible to ANV, because they generally receive more aggressive chemotherapy and have poorer emesis control than older patients.⁵³ Behavioral therapy has been used in patients with ANV.⁵⁴ See the NCCN Guidelines for Antiemesis.

Hematopoietic stem cell transplant

HSCT is a potentially curative therapeutic option for an increasing number of AYA patients with leukemias and lymphomas.⁵⁵ Gonadal dysfunction in males and females related to high-dose conditioning chemotherapy and RT, graft-versus-host disease (GVHD), and chronic immunosuppression are the major post-transplant complications associated with HSCT.^{43, 44} Survivors are also at increased risk for late complications of treatment, which include recurrent infections, secondary cancers, cardiac dysfunction, growth failure, neurocognitive delay, and other end-organ dysfunction.^{43, 44} HSCT survivors are at an increased risk of developing severe or life-threatening chronic health conditions, endocrine complications, or secondary neoplasms compared with non-cancer populations and conventionally treated

cancer patients.⁵⁵ Allogeneic transplant survivors irradiated at 30 years or younger were at higher risk of developing secondary solid cancers.⁵⁶ These findings highlight the increasingly recognized need for long-term follow-up care that incorporates screening and surveillance of AYA survivors of HSCT. See the section on “Survivorship issues: Late effects in AYA cancer survivors.”

Impact of treatment on fertility and fertility preservation

Although preservation of fertility is an issue of crucial importance in AYA patients, it is currently one of the most under prescribed and least implemented services in AYA patients with cancer.^{46, 57} The 2006 American Society of Clinical Oncology (ASCO) Guidelines recommend that providers discuss the options for fertility preservation with all new cancer patients at the time of diagnosis.⁵⁸

Infertility is a major consequence of cancer therapy in both males and females.⁵⁹ The impact of cancer therapy on fertility is related to the age of the patient at the time of treatment and it is dependent on the duration, dose intensity, and type of treatment. Alkylating agent-based chemotherapy is more harmful to the ovaries and testis than chemotherapy regimens containing non-alkylating agents.^{57, 59} High doses of cranial RT can impair hypothalamic pituitary function, resulting in the deficiency of gonadotropin-releasing hormone (GnRH) and impairment in fertility in both males and females.⁶⁰ Gonadal exposure to low doses of radiation can cause oligospermia or azoospermia in males. Higher doses of radiation are associated with both ovarian and uterine dysfunction in women.

NCCN recommendations for fertility preservation

Available evidence strongly supports that fertility preservation is of great importance in AYA patients with cancer and should be an essential part in the management of their cancer.^{46, 57, 61} Women are at



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

risk for premature ovarian failure due to chemotherapy and men are at risk for azoospermia following therapy, which may or may not resolve over time. Fertile Hope has developed a risk calculator based on a compilation of clinical experience and published research on common cancer treatments that may impact reproductive function in both males and females (<http://www.fertilehope.org/tool-bar/risk-calculator.cfm>).

The guidelines recommend that the risk of infertility due to cancer therapy be discussed with all patients at the time of diagnosis and that the provider initiate referral for fertility preservation clinics within 24 hours for appropriate and interested patients.

Fertility preservation for females

Much of the data on the impact of cancer on fertility is from the Childhood Cancer Survivor Study (CCSS) in patients younger than 21 years at the time of diagnosis.^{62, 63} Hypothalamic or pituitary radiation, pelvic RT, and/or increasing alkylating agent doses have been associated with acute ovarian failure and premature menopause.^{64, 65} Total body irradiation (TBI), abdominal and pelvic RT have been shown to cause uterine dysfunction.^{57, 66} Gonadal failure has also been reported among women diagnosed with cancer in their adolescence and young adulthood.⁶⁷ The incidence of gonadal failure is dependent on age at the time of diagnosis and the cumulative dose of alkylating agents.

Fertility is a major concern for young women receiving chemotherapy for breast cancer and HL. Among young women treated with adjuvant chemotherapy for breast cancer, the risk for chemotherapy-related amenorrhea and premature menopause is significantly higher for women with newly diagnosed breast cancer treated with chemotherapy who are over the age of 35.^{68, 69} In a cohort study of 518 female survivors of HL diagnosed between 14-40 years of age, women who were older (22-39 years) at first treatment were at a higher risk for

developing premature menopause after treatment compared to younger patients (14-21 years).⁷⁰ Similarly, the risk of developing premature ovarian failure is also higher among young women, irrespective of the age at the time of treatment (38% for those diagnosed between 30-40 years of age; 37% for those diagnosed between 9-29 years of age) with chemotherapy and RT for HL.⁷¹

Oophoropexy and embryo cryopreservation after in vitro fertilization (IVF) are the two established options for fertility preservation in females.

Oophoropexy involves surgically displacing the ovaries out of the radiation field to minimize ovarian damage and has been shown to preserve ovarian function.⁷² Oophoropexy should be considered for all female patients who will be receiving RT. Ovaries may be surgically moved away from the planned radiation field, either during cancer surgery or in a separate procedure.

For females, if it is possible to delay therapy long enough for a cycle of oocyte stimulation (especially for patients with low- and intermediate-risk HL and low-grade sarcomas), the possibility of embryo cryopreservation should be discussed. Embryo cryopreservation after IVF has been highly successful in women under the age of 40 years.^{46, 57} However, this method requires a male partner or sperm donor.

Mature oocyte cryopreservation and ovarian tissue grafting and freezing are emerging techniques for fertility preservation in young women. They are still considered investigational and their efficacy is unclear. These options are available in some areas, most frequently in the context of clinical trials. Mature oocyte cryopreservation is a potential alternative for single women, but, like embryo



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

cryopreservation, requires hormone stimulation.^{46, 57} Ovarian tissue grafting does not require hormonal stimulation, so there is no long delay in treatment.⁴⁶ However, for some women (eg, those with a malignancy where reimplantation of malignant cells could occur with grafting) this procedure would not be considered appropriate.

GnRH agonists have been used as ovarian protectors during chemotherapy. While some investigators have reported that GnRH agonist administration before and during combination chemotherapy may preserve post-treatment ovarian function in women with breast cancer younger than 40 years,⁷³ others have observed no protection of the ovarian reserve in young women with advanced-stage HL treated with GnRH and escalated BEACOPP (bleomycin, etoposide, doxorubicin, cyclophosphamide, vincristine, procarbazine, and prednisone) chemotherapy.⁷⁴ A more recent systematic review and meta-analysis suggests that although GnRH with chemotherapy in premenopausal women is associated with higher rates of spontaneous resumption of menses and ovulation, it is not associated with improvement in pregnancy rates.⁷⁵ Additional studies are required to confirm these findings. Menstrual suppression does not protect the ovaries. Medroxyprogesterone or oral contraceptives may be used in protocols that are predicted to cause prolonged thrombocytopenia and thus present a risk for menorrhagia.

Management of cancer during pregnancy

The diagnosis of cancer during pregnancy can be an extremely difficult situation for the patient, family and the physician. The most common cancers diagnosed during pregnancy are breast, cervical, ovarian and thyroid cancers, leukemia, lymphoma and melanoma.⁷⁶ Given the rarity of this clinical situation, there is very limited data available on the management of cancer during pregnancy.^{77, 78} AYA women diagnosed with cancer during pregnancy require individualized treatment from a

multidisciplinary team including medical, surgical and radiation oncologists, obstetrician, pathologist and radiologist. Potential benefits and risks of chemotherapy and RT for both the mother and the fetus must be carefully evaluated before the start of treatment. See NCCN Guidelines for Breast Cancer for the management of women with breast cancer during pregnancy. Referral to a gynecologic oncologist and perinatologist with expertise and knowledge of the physiological changes that occur during pregnancy is strongly recommended.

Fertility preservation for males

In males who undergo either chemotherapy or RT, germ cell dysfunction is more common than treatment-induced Leydig cell dysfunction.⁷⁹ Leydig cell dysfunction occurs at RT doses higher than those associated with germ cell dysfunction. AYA men treated with a testicular radiation dose of 20 Gy or more are at high risk for Leydig cell dysfunction, whereas testicular radiation doses of 2 Gy or more can impair spermatogenesis resulting in permanent azoospermia.⁷⁹ TBI used as part of high-dose conditioning therapy prior to HSCT can also affect the testis, resulting in permanent infertility in majority of AYA men undergoing this treatment.⁴⁶ Among patients treated with alkylating agent-based chemotherapy, cumulative doses of cyclophosphamide (20 g/m² or higher) or ifosfamide (higher than 60 g/m²) are associated with a high probability of oligospermia, azoospermia, and infertility, whereas many individuals treated with a cumulative dose of 7.5-10 g/m² or less retain normal sperm production.⁸⁰

Semen cryopreservation and transplantation of spermatogonia are the options for fertility preservation in male patients.^{46, 57} Semen cryopreservation before the start of treatment is the most reliable and well-established means of preserving fertility in AYA males with cancer. Sperm banking should be offered for all patients at the time of diagnosis. AYAs can use either the local sperm bank or the unique



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

collection and preservation kit that is available through Live:On kit (<http://www.liveonkit.com>). Because the age and comfort level of the patients must be taken into account when discussing sperm banking, oncology centers that treat AYA patients should develop a system for offering sperm banking to all AYA patients in a systematic and patient-centered manner. The success of sperm banking may be limited by the fact that some newly diagnosed male cancer patients, such as those with HL, may already have azoospermia associated with the disease.

There is limited evidence regarding the efficacy of hormone suppression in reducing the risk of infertility during chemotherapy.⁸¹ Cryopreservation and subsequent transplantation of spermatogonial stem cells are still experimental and may be an ultimate alternative for some patients in whom semen cryopreservation is not possible.

Psychosocial issues

AYA individuals diagnosed with and treated for cancer have psychosocial issues that are distinct from that of pediatric and adult patients.^{82,83} Some of the challenges faced by AYA patients and survivors include maintaining an active and independent life, coping with treatment-related side effects and stress, seeking and understanding information, accepting cancer, and maintaining a positive attitude.⁸⁴ AYA individuals undergo developmental stages marked by rapid changes in cognitive and emotional growth, and these issues need to be considered to deliver developmentally appropriate psychosocial and supportive care to this population.⁸⁵

Psychosocial needs for AYAs with cancer should be assessed across the following domains: 1) individual function (developmental, emotional, and behavioral issues); 2) relationships (family, peer, and health care

professional); 3) socioeconomic issues; and 4) supportive care services/interventions.

Individual function

Developmental issues

AYAs with cancer have to cope with cancer treatment while accomplishing key developmental tasks such as identity development, including sexual identity; peer involvement; initiating intimate and emotional relationships; establishing autonomy from parents; and independently making decisions about their future that involve education, career, or employment.^{86,87} The impact of diagnosis and treatment of cancer on their physical appearance, sexual development, and sexual function can lead to shame, social isolation, and regressive behaviors if not addressed promptly. Cancer and its often intensive and lengthy treatments put AYA patients at risk for disruptions in their normal activities. Interruptions of school or work due to treatment will have negative consequences for their long-term career opportunities, financial status, and lifetime earnings.⁸⁴ During the treatment period, AYAs should have the opportunity to live as normal a life as possible, continue their education and/or careers, and participate in the many milestones of their lives.⁸⁸

Emotional issues

Cancer-related issues such as confrontation with mortality and loss of fertility can result in significant emotional distress and psychiatric symptoms such as depression and anxiety in AYA patients. These feelings are related to patients' cognitive capacity to understand the severity of their disease while sometimes lacking fully matured cognitive and emotional coping abilities.⁸⁴ Recent studies suggest that the rates of psychological distress are significantly greater amongst AYAs compared to older adults.⁸⁹⁻⁹¹ Kazak et al. recently reported that intensive cancer treatments during adolescence are associated with



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

inferior psychosocial outcomes and health beliefs in survivors compared to their age-matched peers.⁹² Psychological problems are also associated with an increased risk for obesity and poor health behavior, which may increase future risk for chronic health conditions and secondary neoplasms.⁹³

Behavioral issues

AYA patients with cancer may also engage in risky behaviors (tobacco, alcohol, or substance abuse) that may impair their health. Older age at cancer diagnosis, lower household income, less education, no pulmonary-related cancer treatment, and no brain radiation were independently associated with a statistically significant relative risk of smoking initiation.⁹⁴ The risk factors associated with heavy drinking included fair or poor self-assessed health, depression, anxiety, somatization, activity limitations, and cancer-related fears and uncertainty.⁹⁵ Low perception of susceptibility to late effects, older adolescence compared to early adolescence, and worry were the strongest predictors of substance abuse.⁹⁶

While AYA patients may be aware of the complications associated with tobacco, alcohol, or substance abuse during their treatment, they may not avoid them throughout their treatment, as these habits make them feel normal and like part of their peer group. Clinicians working with this population need to be aware of this and address the issues in a sensitive and confidential manner.⁸⁸

NCCN recommendations for supportive care services/interventions

- For all AYA patients, provide counseling regarding the risks of treatment-related infertility and discuss options for fertility preservation prior to the start of therapy.⁹⁷
- Provide AYA patients with flexible treatment dates, consultation times, and procedures to enable them to continue with their

treatment without interrupting their school/work or other normal activities.⁸⁸

- Offer psychosocial support and counseling to help alleviate distress. See the NCCN Guidelines for Distress Management.
- Refer AYA patients with cognitive dysfunction or other psychiatric symptoms (eg, depression or anxiety) to a mental health provider and community-based resources serving AYA patients.
- Refer patients with signs, symptoms, and a history of substance abuse or addiction to a risk reduction or substance abuse management program.
- Since the incidence of sexually transmitted infections peaks among AYAs 15-24 years of age, provide preventative health education about sexually transmitted diseases.⁸⁸
- Recommend HPV immunization (if not previously administered) for the prevention of secondary cancers since the vaccine has been shown to prevent cervical carcinoma and anal epithelial neoplasia, the precursor to carcinoma.⁹⁸
- Prescribe and provide nutrition and exercise recommendations for all AYAs.
- Refer patients experiencing challenges with their faith or belief in a just or fair world to faith-based resources or activities (eg, church youth groups, mentors).⁸⁴

Adherence to treatment

Adherence is defined as the extent to which a person's behavior corresponds with agreed recommendations from a health care provider. Nonadherence to recommended treatment and follow-up care contribute to poor clinical outcomes in AYAs with cancer.^{99, 100} Failure to keep up with appointments can lead to delayed identification of side effects, complications, or secondary cancers.



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

Nonadherence to treatment regimens has been an ongoing problem among patients with cancer, and the prevalence of nonadherence has been consistently higher among adolescents compared to younger or older patients with cancer.⁹⁹ Nonadherence to oral chemotherapy contributes to reduced treatment efficacy and increased risk of recurrence. Available evidence from clinical trials that have included AYA patients with leukemia and lymphoma suggests that a substantial portion of AYA patients with cancer (27% to 63%) have difficulties adhering to their oral treatment regimens.^{99, 100}

Nonadherence to other components of cancer treatment (eg, failure to keep appointments for treatment or follow-up, refusing medical examinations, preparing for procedures or therapy) was also identified in AYA patients. Treatment nonadherence in clinical trials can interfere with adequate evaluation of the efficacy of a given treatment regimen, which in turn can invalidate the results of a clinical trial.

Risk factors for nonadherence among AYA patients with cancer include patients' emotional functioning (depression and self-esteem), personal beliefs (perceived severity of cancer diagnosis and the necessity of intervention), growing independence, competing obligations (school, work, and family), and lack of insurance and appropriate psychosocial support.¹⁰¹ In a randomized controlled trial, video game intervention significantly improved treatment adherence to prophylactic antibiotics among adolescents and young adults with acute leukemia, lymphoma, and soft tissue sarcoma.¹⁰² A recent meta-analysis showed that behavioral and multicomponent interventions have been shown to have a moderate effect on improving treatment adherence in children (2 to 15 years) with chronic conditions such as diabetes, asthma, and cystic fibrosis.¹⁰³

Additional studies evaluating the effect of interventions to improve adherence in AYA patients with cancer are needed. In the absence of data from studies evaluating the effect of interventions to improve adherence in AYA patients with cancer, the findings from the studies involving AYA patients with other chronic diseases could be extrapolated to this patient population.

NCCN recommendations to promote adherence

- Provide education and/or guidance about each medication prior to the start of treatment and every time there is a change in treatment. Review the list of medications as well as their dose, purpose, and adverse effects.^{99, 100}
- Modify treatment protocol (eg, simplify dosing schedule, change timing and frequency of medication or method of administration), when medically possible, to fit into an AYA's lifestyle and normal activities.^{99, 100}
- Provide access to systematic and standardized symptom management for side effects related to cancer treatment.^{99, 100} See the NCCN Guidelines for Supportive Care.

Relationships

Social, peer, and family relationships

AYAs often have to endure lengthy hospital stays under the supervision of health care providers, resulting in significant isolation from their family members and peer group.⁸⁸ Isolation and alienation are common among AYA individuals diagnosed with cancer, because they often miss out on the life experiences shared by their non-ill peers. Reinforcing relationships with family, peers, and health professionals is an important aspect of life for AYAs with cancer.^{85, 104}

While some studies have identified family support and cohesiveness as important contributors to a survivor's adjustment, others have identified



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

the important role played by same-aged peers (healthy peers as well as other AYA cancer survivors) in helping AYAs cope with cancer and overcome feelings of loneliness.^{82, 85} In one study, AYAs with cancer (16-22 years) identified social support (friends and health care providers) as their major coping strategy to deal with cancer, whereas family support was identified as their important source for emotional support.¹⁰⁵ In another study, AYA patients and survivors reported that opportunities to meet other young adult survivors were more important than the support they received from family and peers.⁹⁷

Peer support programs assist AYA patients and survivors in establishing and maintaining relationships with their normal peers as well as with other AYAs with cancer, offer opportunities to achieve age-related developmental tasks (building interpersonal and problem-solving skills), and promote positive psychosocial growth.^{97, 106} Peer support also provides AYAs with an opportunity to address some of their concerns, such as coping with uncertainty about the future, establishing autonomy while being increasingly dependent on family and friends, sexual identity, and infertility, thereby reducing feelings of social isolation.¹⁰⁶

AYA peer support groups have been developed in a variety of formats, including face-to-face meetings, camp style formats, or online support groups.¹⁰⁷ Summer camps and adventure programs where participants are physically challenged have resulted in improvements in self-confidence, independence, and social contacts.^{84, 107} Many of the AYA patients may not be interested in conventional cancer support groups but are willing to participate in social networking events involving other AYA patients, survivors, and family members.⁸⁴ Indeed, studies of AYA patients and survivors indicated that 73% of patients currently receiving therapy and 74% of off-treatment survivors reported that their needs for retreats and camp programs were unmet.^{108, 109}

Communications with health care professionals

Communicating information to AYAs can be challenging, especially since there are several subgroups within the AYA population with different levels of cognitive and emotional development. It is very important to establish direct communication with the patients on an individual basis, with sufficient sensitivity to each patient's needs and preferences.²⁴ While some patients prefer not to receive direct communication about their cancer, others may desire to take a more prominent role in the management of their care. For the latter group, information should be provided directly to the patient in an age-appropriate manner, allowing time to process the information and deliver information in a caring manner.¹¹⁰ AYAs prefer that information about their cancer and cancer-related risks be communicated to them in a manner that is positive, respectful, and nonjudgmental.⁸⁸ In a pilot project aimed at eliciting the views of AYA patients with cancer, humor, closely followed by expertise and knowledge, was identified as the most important characteristic that patients would like to see in their nurses.¹¹¹ Since there is evidence that AYA patients are willing to use the internet to get health information and support, it will also be helpful to provide them with a list of recommended and reliable age-appropriate online sources to access information about their cancer, particularly with regard to treatment and late effects, fertility preservation, mental health counseling, peer support groups, diet, and nutrition.^{107, 109, 112} See Online Resources for AYA patients and Survivors in the guidelines.

NCCN recommendations for supportive care services/interventions

- Promote communication between AYA patients and family members (parents, spouse/partners, and siblings).¹¹
- Provide information to family members and partners about psychosocial support and behavioral services to increase



awareness of the possible psychosocial issues associated with diagnosis of cancer in AYAs.

- Consider family-based intervention models from pediatrics (eg, parent support groups, Impact of Traumatic Stressors Interview Schedule).⁸
- Establish direct communication with the individual patients, providing age-appropriate information about their cancer, treatment options, and potential side effects, thus reinforcing the importance of AYA involvement in decision-making.^{24, 97}
- Some AYA patients prefer not to share information about their cancer with their family in an effort to shield their family members from some of the things they themselves worry about. Therefore, obtain their permission to share information with other family members.
- Provide information about peer support groups and create flexible visiting hours and an environment that will encourage peers to visit AYA patients.⁸⁸

Socioeconomic issues

AYAs are much more likely to be uninsured or underinsured individuals than adults or children, with many of them in a transition between their parents' insurance and their independent insurance.⁸ Young adult survivors of childhood cancers are more likely to report health-related unemployment, lower rates of health insurance coverage, and more difficulties obtaining coverage compared to their siblings.¹¹³

Furthermore, unemployment and lack of health insurance appear to be significant predictors of psychological distress in the childhood cancer survivor population.¹¹⁴ Uninsured AYA patients are also less likely to participate in clinical trials.³⁸ As described above, advanced stage of cancer at diagnosis and lack of health insurance were significantly associated with longer time to cancer diagnosis in AYAs.²³ Greater

rates of unemployment and lack of health insurance among AYA patients and survivors are also associated with limited access to long-term follow-up care.⁸⁴ AYA patients with employment also experience problems in obtaining health and life insurance due to their pre-existing cancer history.⁸² Even those with relatively comprehensive insurance may be liable for substantial out-of-pocket expenses related to treatment, such as transportation costs associated with traveling for treatment, accommodations, meals, and childcare as well as non-treatment-related costs.⁸² Financially independent AYA patients also have to face an additional burden of loss of income because of their inability to work during treatment. Once the treatment is over, AYA patients with cancer also need long-term follow-up care for monitoring and treatment of late effects.

NCCN recommendations for supportive care services/interventions

- Assess AYA patients' health insurance status and potential and provide information on potential sources of coverage (eg, Medicaid, Social Security, and Disability Insurance) and other key elements associated with insurance coverage.
- Educate AYA patients about the benefits for which they may qualify (eg, short or long-term disability, state disability benefits, Social Security benefits, food stamps).
- Provide a referral for transportation assistance programs (eg, van ride programs, voucher programs) for AYA patients who have to travel to receive treatment. Identify resources for respite care that would be helpful for those with young children.
- For those who desire to receive complementary and alternative medicine (CAM), refer them to reputable providers of CAM services.
- Provide information about reliable online sources to access age-appropriate information related to their cancer. See Online Resources for AYA patients and survivors in the guidelines.



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

- Educate AYA patients with cancer about their long-term follow-up care for monitoring and treatment of late effects, long after completion of treatment.
- Integrate financial assistance for AYA cancer survivors into their survivorship plans.

Survivorship issues

Late effects in AYA cancer survivors

AYA cancer survivors are at increased risk for late effects related to cancer treatment, and the risk for long-term effects is dependent on the age at initial diagnosis and the type of treatment.¹¹⁵⁻¹¹⁷ In addition, the risk for many late effects may also be influenced by family history, lifestyle behaviors, and comorbid health conditions. Much of the understanding of the long-term outcomes of AYA cancer survivors comes from the CCSS, which includes long-term survivors of childhood and adolescent cancers who were diagnosed prior to age 21.^{118, 119} No such large cohort studies have addressed the survivorship issues related to cancer diagnosed in young adult patients between the ages of 22 and 39 years. Among adult survivors of childhood and adolescent cancer, Oeffinger et al. reported that by 30 years after the cancer diagnosis, the cumulative incidence of a chronic health condition was 73%, with a cumulative incidence of 42% for severe, disabling, or life-threatening conditions or death. Importantly, the risk for a chronic health condition (ie, long-term or late effect) was similar for those diagnosed with the primary cancer in adolescence and in childhood.¹¹⁸ Age at treatment exposure modifies the risk of some late effects (eg, breast cancer following chest radiation, cardiomyopathy following anthracycline chemotherapy) but not others (eg, ischemic coronary artery disease following chest radiation).^{120, 121}

While several single cancer studies have assessed long-term outcomes among HL and testicular cancer survivors across the AYA age range, the long-term outcomes of survivors of other cancers occurring in young adulthood, such as breast, ovarian, and thyroid cancers or melanoma, remain understudied. Outcomes from the CCSS among those diagnosed between the ages of 15 and 20 are particularly relevant for the NCCN Guidelines for AYA Oncology. Since there is a paucity of literature on survivorship issues related to cancer diagnosed during adolescence and young adulthood, the findings from the CCSS and similar studies focusing on childhood and adolescent cancer survivors could be extrapolated to the survivors of AYA cancers, albeit with caution.

Some of the more common late effects among AYA cancer survivors are discussed below.

Secondary cancers

AYA cancer survivors (diagnosed between 15-39 years) are at significant risk of developing a variety of secondary cancers compared to the general population.¹²² The risk and specific types of secondary cancers are widely dependent on the type of initial cancer diagnosis and treatment exposure.¹²³⁻¹²⁵ Older age at diagnosis (15-21 years) was associated with increased risk for breast cancer, nonmelanoma skin cancers, and other solid organ cancers (including head and neck, small intestine, and colorectal cancers).^{125, 126}

AYA survivors of HL diagnosed between 21-39 years of age are at increased risk of developing secondary cancers.¹²⁴ The most frequently observed secondary cancers are breast, lung, thyroid, and gastrointestinal cancers.¹²⁰ Adolescent or young women treated with chest radiation for HL are at significantly increased risk of developing secondary breast cancer, and the risk for secondary breast cancer



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

among HL survivors is strongly associated with age at diagnosis and mediastinal radiation dose.¹²⁷⁻¹³⁰ In a cohort of 770 female survivors who had been diagnosed with HL before age 41 years, the risk of developing breast cancer increased with increasing radiation dose (38.5 Gy or more).¹²⁸ In an international, population-based study of 3,817 female survivors of HL diagnosed at age 30 years or younger, Travis et al. reported that for women treated at age 25 years with a chest radiation dose of at least 40 Gy without alkylating agents, the estimated cumulative absolute risks of developing breast cancer by age 35, 45, and 55 years were 1.4%, 11.1%, and 29.0%, respectively.¹²⁹

Alkylating agent-based chemotherapy for HL has been associated with a modestly increased risk for secondary lung cancers in patients diagnosed at 40 years or younger, and the risk increased with both increasing number of cycles of alkylating agents and the cumulative dose.¹³¹ In this study, the risk of secondary lung cancer was substantially higher among survivors who smoked (9.6% due to treatment alone compared to 63.3% due to the combination of treatment and smoking). In a recent collaborative British Cohort study that assessed the risk of developing secondary cancers in 5,798 patients diagnosed with HL between 15-34 years of age, the 20-year cumulative risk of second cancer was 13% and 18%, respectively, for chemotherapy alone and combined modality therapy.¹³² Risks for secondary lung cancer, NHL, and leukemia were significantly higher after treatment with chemotherapy alone, whereas combined modality therapy was associated with a higher risk for these and several other cancers.¹³²

AYA survivors of testicular cancer are also at significantly increased risk of developing secondary cancers, including contralateral testicular cancer, leukemia, malignant mesothelioma, and cancers of the lung, colon, esophagus, stomach, and pancreas.^{133, 134} In a population-based

study of 29,515 testicular cancer survivors, the 15-year cumulative risk of developing contralateral testicular cancer was almost 2%, which is 12-fold higher than that of the general population.¹³⁵ In an international, population-based study of 40,576 testicular cancer survivors, the cumulative risk of developing solid tumors by age 75 years was slightly higher for seminoma patients than for nonseminoma patients diagnosed at 35 years of age (36% and 31%, respectively).¹³⁶ The combination of chemotherapy and RT was associated with a larger risk of secondary solid tumors than RT alone, although the difference was not statistically significant.¹³⁶ Secondary leukemia related to chemotherapy with topoisomerase II inhibitors and alkylating agents has also been reported in testicular cancer survivors. In one study, the cumulative incidence of secondary AML was 0.5% at 2 years after treatment with high-dose chemotherapy (with a median cumulative etoposide dose of 4.9 g/m²) and autologous stem cell transplantation.¹³⁷ In another study involving 42,722 one-year survivors of testicular cancer, the estimated excess cumulative leukemia risk was 0.23% at 30 years after testicular cancer diagnosis.¹³⁸ The risk for secondary AML was higher for patients treated with chemotherapy compared to those treated with radiotherapy alone.

The risk for secondary malignancies among survivors of cervical and breast cancers, NHL, and melanoma has been assessed in only a few cohort studies.¹³⁹⁻¹⁴¹ Among 104,760 one-year survivors of cervical cancer, patients heavily treated with RT were at increased risk for second cancers at sites in close proximity to the cervix beyond 40 years of follow-up. The 40-year cumulative risk for any second cancer was higher among women diagnosed before age 50 than among women diagnosed after age 50 (22.2% and 16.4%, respectively).¹³⁹ In a population-based cohort of 376,825 one-year survivors of breast cancer from the Scandinavian cancer registries, women diagnosed at 40 years

or younger with localized disease were particularly at risk of developing a second cancer at 30 or more years after breast cancer diagnosis.¹⁴⁰ In an analysis of 28,131 patients from the Swedish Cancer Registry, the risk of developing subsequent solid tumors after NHL during the first decade was higher among patients diagnosed between 20-39 years of age compared to those who were age 40 years or older at the time of diagnosis.¹⁴¹ In the SEER database analysis of 89,515 melanoma survivors, patients diagnosed at younger than 30 years of age had the highest risk of developing secondary cancers (breast, prostate, and NHL being the most common cancers) at more than 20 years after initial diagnosis.

Long-term AYA survivors of pediatric-predominant cancers, including ALL, CNS tumors, and bone and soft tissue sarcomas, are also at risk of developing secondary cancers. The risk is especially higher among patients diagnosed at a younger age (17 years or younger for ALL and CNS tumors; 18 years or younger for bone and soft tissue sarcomas).¹²² Among long-term survivors of bone cancers at 25 years after diagnosis, the cumulative incidence of subsequent cancers is higher for those diagnosed with Ewing's sarcoma compared to those diagnosed with osteosarcoma (9.0% and 5.4%, respectively).^{142, 143}

Clinicians who provide care for the majority of AYA cancer survivors must implement and evaluate methods for improving awareness of secondary cancers. They must also implement appropriate surveillance strategies for early detection of these malignancies.¹⁴⁴ An annual breast magnetic resonance imaging (MRI) and mammogram are recommended for women treated with a chest radiation dose of 20 Gy or more prior to age 30 years. A colonoscopy is recommended starting at age 35 or 10 years after radiation, whichever occurs last, for patients treated with abdominal or pelvic radiation of 30 Gy or more.

Cardiovascular complications

Cardiovascular complications (congestive heart failure [CHF], myocardial infarction [MI], pericardial disease, and valvular abnormalities) are the leading non-malignant cause of death among survivors of AYA cancers, compared to the general population.¹⁴⁵ Mediastinal irradiation and anthracycline-based chemotherapy are the strongest risk factors for late cardiovascular complications in AYA survivors of HL.^{121, 146, 147} In the British Cohort study of 7,033 HL patients, the risk of death from MI was highest for patients younger than 35 years at the time of treatment with supradiaphragmatic RT.¹⁴⁷ Patients treated with anthracyclines were at increased risk for MI within one year after first treatment, whereas the risk for MI among patients treated with supradiaphragmatic RT and vincristine without anthracyclines increased sharply after the first year of follow-up.¹⁴⁷ In another study of 1,474 survivors of HL younger than 41 years at the time of treatment, mediastinal RT increased the risk of MI, CHF, and valvular disorders, whereas the addition of anthracyclines to RT elevated the risks of CHF and valvular disorders.¹²¹ The 25-year cumulative incidence of CHF after mediastinal RT and anthracyclines was 8%.

Cisplatin-based chemotherapy is associated with long-term risk for cardiovascular complications in testicular cancer survivors.^{148, 149 150} In a Dutch study of 2,512 testicular cancer survivors, nonseminoma testicular cancer survivors younger than 30 years at diagnosis treated with mediastinal irradiation and chemotherapy with cisplatin, vinblastine, and bleomycin were at increased risk for MI within 20 years of treatment.¹⁴⁸ More recently, Haugnes et al. reported that treatment with cisplatin, bleomycin, and etoposide and/or RT was associated with increased risks for cardiovascular disease in testicular cancer survivors; chemotherapy alone or in combination with RT significantly increased the risk for MI.¹⁴⁹



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

Survivors of brain tumors, leukemia, NHL, and bone and soft tissue sarcomas treated with anthracyclines and cardiac irradiation are also at significantly higher risk of adverse cardiovascular complications. However, the majority of patients included in these studies were younger than 21 years at the time of diagnosis.¹⁵¹

Pulmonary complications

Chemotherapy and chest radiation are associated with pulmonary toxicity and can compromise pulmonary function in survivors of AYA cancer.^{150, 152} Age at diagnosis (15-21 years compared with less than 15 years) and pulmonary toxic chemotherapy alone or combined with chest radiation were associated with a significantly increased relative risk of lung fibrosis and pleurisy.¹⁵² The cumulative incidence increased up to 15-20 years after diagnosis. Other complications include recurrent pneumonia, chronic cough, supplemental oxygen use, and shortness of breath.

A large international study reported a significant increase in mortality from respiratory diseases among testicular cancer survivors treated with chemotherapy compared to the general population.¹⁵³ Risk factors for pulmonary toxicity include age at diagnosis, cumulative bleomycin dose, reduced glomerular filtration rate, renal dysfunction, and stage IV disease at presentation.¹⁵⁴ Recently, Haugnes et al. reported that among 1,049 testicular cancer survivors, those treated with large cumulative cisplatin doses or chemotherapy combined with pulmonary surgery had significantly reduced pulmonary function compared with those treated with surgery alone.¹⁵⁵ Bleomycin dose was not associated with restrictive lung disease. Instead, in a multivariate model, cisplatin dose ($p = 0.007$) and age at diagnosis ($p = 0.008$) were associated with the risk for restrictive lung disease.

Neurological complications

AYA survivors of brain tumors treated with cranial RT are at increased risk for neurologic complications, including hearing impairments, cataracts and other vision problems, seizure disorders, and coordination and motor control problems.^{156, 157} However, these findings are relevant to survivors diagnosed at 21 years of age or younger.

Long-term AYA survivors of testicular cancer who were treated with cisplatin-based chemotherapy are at risk for neurological complications such as sensory neuropathy, tinnitus, hearing impairment, and Raynaud's phenomena (white or cold hands or feet on cold exposure).¹⁵⁰ Among 1,814 survivors of testicular cancer included in a Norwegian observational study, Raynaud-like phenomena were the most frequently reported complications (39% of men), followed by paresthesia of the hands or feet (29%), and tinnitus and hearing impairment (22% and 21%, respectively) by men treated with chemotherapy compared to those not treated with chemotherapy.¹⁵⁸ The incidences of paresthesia of the feet were also higher among men treated with RT.

Stroke, although relatively uncommon, is a devastating neurological complication in AYA survivors of brain tumors and leukemia treated with cranial RT and HL treated with mantle field radiation.^{159, 160} In a retrospective cohort study of 2,201 5-year survivors of HL, those treated with RT to the neck and mediastinum were particularly at increased risk for stroke and transient ischemic attack.¹⁶¹ The incidences were higher among patients diagnosed at younger than 21 years than those diagnosed between 21 and 30 years of age. The standardized incidence ratio was 3.8 and 3.1 respectively.

Nephrotoxicity

Long-term renal dysfunction has been reported in survivors of testicular cancer treated with infradiaphragmatic RT and cisplatin-based chemotherapy. In one study with a long-term follow-up, renal impairment was observed in 8% of patients treated with abdominal RT alone compared to a 14% reduction in patients with chemotherapy with or without RT. Age at treatment and type of treatment were associated with impaired renal function.

Endocrine complications

Cranial or spinal RT, TBI, and target organ irradiation involving the neck, abdomen, pelvis, and testes are associated with endocrine late effects in survivors of AYA cancers.⁷⁹ The most common endocrine complications include growth hormone (GH) deficiency, thyroid gland abnormalities, gonadal dysfunction, and decreased fertility.^{43, 60} AYA cancer survivors treated with RT dose of 18 Gy or more to the hypothalamic-pituitary-adrenal (HPA) axis are at high risk for GH deficiency, whereas those treated with RT dose of 40 Gy or more to the HPA axis are at risk for developing central hypothyroidism, gonadotropin deficiency, and central adrenal insufficiency.

GH deficiency can be observed within 5 years after treatment with RT doses higher than 30 Gy, whereas in patients treated with lower doses (18-24 Gy) it may not be evident for 10 years or more.⁷⁹ Secondary thyroid cancers, hypothyroidism, and, to a lesser extent, hyperthyroidism are more common among AYA survivors of brain tumors, ALL, HL, and those who underwent HSCT.^{156, 162, 163} Testicular cancer survivors treated with chemotherapy and RT are at greater risk for hypogonadism.¹⁵⁰ Low testosterone levels and testosterone replacement have been reported in 34% and 4% of testicular cancer survivors, respectively.¹⁶⁴

Long-term follow-up

As discussed above, AYA cancer survivors have a high risk of developing a wide range of late effects. AYA cancer survivors may benefit from regular screening and early intervention for cardiovascular disease.¹⁶⁵ Continued follow-up of AYA cancer survivors is needed to monitor the pulmonary complications.¹⁶⁶ Development of a “Cancer Treatment Summary and Survivorship Care Plan,” including periodic evaluation with focused history, physical examination, and screening based upon treatment exposures, and risk for treatment-related late effects, should be an integral part of management of AYA patients with cancer.^{116, 167, 168}

The models for AYA survivorship care include cancer center follow-up (primary treatment team or specialized long-term follow-up clinics), follow-up by the patient's primary care physician, or a combination of both (shared care model).^{168, 169} Some studies have shown that a shared care model involving both the primary oncology team and the primary care physician is feasible and can facilitate appropriate care in childhood cancer survivors.¹⁷⁰⁻¹⁷²

Risk stratification of survivors based on the current medical issues and prior treatments may be helpful to determine the different levels of follow-up in the shared care model.^{169, 173, 174} Survivors at low risk for late effects (treated with surgery alone and/or chemotherapy with no radiation, not including alkylating agents, anthracycline, bleomycin, or epipodophyllotoxin) can be transitioned to their primary care physician soon after completion of therapy. Survivors at moderate risk for late effects (treated with low- or moderate-dose chemotherapy with no radiation, containing alkylating agents, anthracycline, bleomycin, or epipodophyllotoxin) can be evaluated by their oncology team or primary care physician on alternating years. Survivors at high risk for late effects, such as those treated for CNS cancers or those treated with a



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

stem cell transplantation, any radiation, high-dose alkylating agents, anthracycline, bleomycin, or epipodophyllotoxin, should be followed annually by their oncology team and continue follow-up care with their primary care physician.

NCCN recommendations

The following screening recommendations are adapted from the Children's Oncology Group (COG) Long-Term Follow-Up Guidelines for Survivors of Childhood, Adolescent, and Young Adult Cancers, available at survivorshipguidelines.org.¹⁴⁴ The recommendations are based upon the treatment exposure; timing and intensity of screening may be adapted based upon additional risk factors.

Cardiovascular screening

- Cardiovascular risk assessment and screening are recommended for AYA cancer survivors treated with mediastinal, chest, or abdominal radiation of 30 Gy or more.
- An echocardiogram (or MUGA scan) every 1-2 years is recommended for those treated with anthracycline-based chemotherapy, chest radiation, or combined anthracycline-based chemotherapy and chest radiation.
- Screening for ischemic coronary artery disease remains controversial; cardiology consultation (5 to 10 years after radiation) in patients who received chest radiation of 40 Gy or more can be considered.

Pulmonary screening

- Chest x-ray and pulmonary function tests as a post-therapy baseline and then as clinically indicated in patients with abnormal results or progressive pulmonary dysfunction is recommended for all patients treated with chest radiation, alone or in combination with

chemotherapy containing bleomycin, busulfan or carmustine, TBI, and bleomycin-based chemotherapy.

Neuroendocrine screening

- Routine neuroendocrine axis screening to monitor GH deficiency, central hypothyroidism, gonadotropin deficiency, and central adrenal deficiency is recommended throughout the entire lifespan of AYA cancer survivors.
- Periodic testing of thyroid function and screening for early detection of thyroid cancer (thyroid stimulating hormone [TSH] and thyroid/neck exam, annually) must be performed on AYA cancer survivors.
- Audiology testing as a post-therapy baseline and then as clinically indicated for signs and symptoms of hearing loss is recommended for patients with cisplatin-based chemotherapy, radiation involving the ear, or a combination of cisplatin-based chemotherapy and cranial/ear radiation.

Neuropsychological evaluation

- Severe neurocognitive deficits are uncommon in survivors of AYA cancer, including CNS tumors. However, subtle deficits in executive function, sustained attention, memory, and processing speed may occur with a cranial radiation dose of 18 Gy or more.
- In patients with evidence of impaired educational or vocational progress, formal neuropsychological evaluation is recommended.

Assessment of renal function

- Screening for renal insufficiency and secondary renal/renovascular hypertension is recommended for all patients treated with a radiation dose of 10 Gy or more and a combination of radiation with chemotherapy containing nephrotoxic drugs such as cisplatin, ifosfamide, aminoglycosides, amphotericin, and immunosuppressants.



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

- Screening for hemorrhagic cystitis/ bladder fibrosis or bladder cancer is recommended for patients treated with cyclophosphamide combined with pelvic irradiation.

Assessment for gonadal function

Males

- Analyze semen for evaluation of infertility or as requested by the patient; periodic evaluation over time is recommended as resumption of spermatogenesis can occur up to 10 years after treatment.
- Monitor testosterone levels as clinically indicated in patients with clinical signs and symptoms of testosterone deficiency.
- See also the section on “Impact of treatment on fertility and fertility preservation.”

Females

- Monitor follicle-stimulating hormone (FSH), luteinizing hormone (LH), estradiol levels as indicated in patients with irregular menses, primary or secondary amenorrhea, and/or clinical signs and symptoms of estrogen deficiency.
- See also the section on “Impact of treatment on fertility and fertility preservation.”

Palliative and End-of-life care

Palliative care is interdisciplinary care of patients with life-threatening illnesses, malignant as well as non-malignant. The goal of palliative care in a patient with incurable cancer is to control symptoms, relieve suffering from adverse effects of treatment, and improve quality of life for patients and their families, regardless of the stage of the disease or the need for other therapies.¹⁷⁵ See NCCN Guidelines for Palliative Care.

Palliative care services for AYA patients should be provided by a multidisciplinary team with expertise in understanding the psychosocial, emotional, developmental, and financial issues that are unique to this age group.¹⁷⁵ Introduction of palliative care for symptom management and psychosocial support should occur before the patient is considered “palliative” in order to provide the best possible care for the patient.¹⁷⁶ Palliative care is appropriate even when patients are being treated with curative intent, and there is growing consensus that AYA patients should have access to palliative care services from the time of diagnosis until the time of death or cure.¹⁷⁶ Patients, caregivers, and health care professionals should be taught that palliative care is an integral part of their comprehensive cancer care. AYA patients usually are not making decisions in isolation. While some AYA patients have the ability to make life and death decisions independently, many are either not the primary decision maker or they rely intensely on input from parents, spouses, significant others, and other family members.¹⁷⁵ Palliative care services should also consider the psychosocial needs of the patient’s family, friends, and caregivers.¹⁷⁶ Social support is required for almost all AYA patients receiving palliative care.

End-of-life care involves the management of delirium, existential distress, discussion about the place of death, and support of family.¹⁷⁵ AYA patients understand that death is permanent and irreversible. It is imperative for health care professionals not to assume that AYA patients may be less inclined to discuss death and other end-of-life issues.¹⁷⁵ In an exploratory study of 50 adolescent patients (15-21 years of age) with and without chronic illnesses, adolescents were willing to discuss end-of-life decision making by taking part in a one-on-one survey administered by a researcher.¹⁷⁷ AYA patients’ opinions about end-of-life care vary across this age group. Exploring



NCCN Guidelines Version 1.2012 Adolescent and Young Adult Oncology

individual preferences for end-of-life care and providing interventions specific to the needs of this patient population could significantly improve end-of-life care.¹⁷⁸ In one retrospective review, a significant number of adolescents dying of cancer felt that discussions about the end-of-life occurred very close to death, thus allowing very little time to psychologically prepare for death.¹⁷⁹ The palliative care team can also help to relieve physical and emotional suffering as well as facilitate difficult end-of-life issues such as nutrition/hydration, sedation, treatment cessation, and place of death.¹⁷⁵ An advance care planning document may be appropriate and helpful for terminally ill AYA patients with metastatic cancer.¹⁸⁰

Table 1. Incidences of cancer (all sites combined) by age group and sex in the AYA population^{a,b}

Age group	Year	No. of cases	
		Male	Female
15-19	2008	2,536	2,176
20-24	2008	3,596	3,753
25-29	2008	5,026	6,785
30-34	2008	6,403	10,857
35-39	2008	9,607	18,895

a. Data is from United States Cancer Statistics: 1999 - 2008 Incidence, WONDER Online Database. United States Department of Health and Human Services, Centers for Disease Control and Prevention and National Cancer Institute; 2011. Accessed at <http://wonder.cdc.gov/cancer-v2008.html> on Feb 19, 2012.

b. Data for in situ Breast Cancers are listed separately from Breast Cancers and not included in the "all sites" category.



Table 2. Age-specific SEER incidences^a of cancer by age group and sex in the AYA population (2004-2008)^b

Cancer type	Ages 15-19		Ages 20-24		Ages 25-29		Ages 30-34		Ages 35-39	
	Male	Female								
Bone sarcomas	2.1	1.3	1.2	0.9	0.7	0.7	0.6	0.6	0.7	0.6
Carcinoma of breast	-	-	-	1.4	-	7.8	-	26.1	0.1	59.5
CNS cancers	2.2	2.1	2.2	2.3	3.1	2.2	3.6	2.6	4.2	3.0
Carcinoma of cervix and uterus	-	-	-	1.7	-	6.3	-	14.3	-	20.9
Carcinoma of colon and rectum	0.3	0.3	0.9	1.0	2.1	1.9	4.4	4.5	9.3	8.3
Carcinoma of respiratory tract	-	-	0.3	0.3	0.5	0.6	1.1	1.3	2.8	3.1
Germ cell neoplasms	4.3	1.1	10.6	1.2	14.0	1.1	13.4	0.9	11.1	0.7
Leukemias	3.7	2.6	3.0	2.3	3.0	2.2	3.4	2.8	4.2	3.2
Lymphomas										
HL	3.0	3.1	4.3	4.7	4.0	4.5	4.0	3.8	3.4	2.7
NHL	2.1	1.5	2.9	1.9	3.8	2.8	5.5	4.0	8.6	5.5
Melanoma	1.2	1.9	2.5	6.0	4.7	9.9	7.4	12.3	10.9	16.5
Soft tissue sarcomas	1.4	1.5	2.0	1.7	3.1	2.1	4.2	2.8	5.9	3.7
Thyroid carcinoma	0.7	3.4	1.3	8.8	2.7	14.2	4.0	20.2	5.3	24.6

a. Rates are per 100,000

b. Data from Howlader N, Noone AM, Krapcho M, Neyman N, Aminou R, Waldron W, Altekruse SF, Kosary CL, Ruhl J, Tatalovich Z, Cho H, Mariotto A, Eisner MP, Lewis DR, Chen HS, Feuer EJ, Cronin KA, Edwards BK (eds). SEER Cancer Statistics Review, 1975-2008, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2008, based on November 2010 SEER data submission, posted to the SEER web site, 2011.



References

1. Bleyer A, O'Leary M, Barr R, Ries L. Cancer Epidemiology in Older Adolescents and Young Adults 15 to 29 Years of Age, Including SEER Incidence and Survival: 1975-2000. National Cancer Institute, NIH Pub. No. 06-5767 2006. Available at: <http://www.seer.cancer.gov/publications/aya/>.
2. Burke ME, Albritton K, Marina N. Challenges in the recruitment of adolescents and young adults to cancer clinical trials. *Cancer* 2007;110:2385-2393. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17918260>.
3. Ferrari A, Montello M, Budd T, Bleyer A. The challenges of clinical trials for adolescents and young adults with cancer. *Pediatr Blood Cancer* 2008;50:1101-1104. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18360838>.
4. Bleyer A, Budd T, Montello M. Adolescents and young adults with cancer: the scope of the problem and criticality of clinical trials. *Cancer* 2006;107:1645-1655. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16906507>.
5. Bleyer A. Young adult oncology: the patients and their survival challenges. *CA Cancer J Clin* 2007;57:242-255. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17626120>.
6. Bleyer A, Barr R, Hayes-Lattin B, et al. The distinctive biology of cancer in adolescents and young adults. *Nat Rev Cancer* 2008;8:288-298. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18354417>.
7. Ramphal R, Meyer R, Schacter B, et al. Active therapy and models of care for adolescents and young adults with cancer. *Cancer* 2011;117:2316-2322. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523752>.
8. Closing the gap: Research and care imperatives for adolescents and young adults with cancer: Report of the adolescent and young adult oncology progress review group: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, Livestrong™ Young Adult Alliance 2006. Available at: http://planning.cancer.gov/library/AYAO_PRG_Report_2006_FINAL.pdf.
9. Bleyer A, Viny A, Barr R. Cancer in 15- to 29-Year-Olds by Primary Site. *Oncologist* 2006;11:590-601. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16794238>.
10. Howlader N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2008, based on November 2010 SEER data submission, posted to the SEER web site, 2011: National Cancer Institute, Bethesda, MD,. Available at: http://seer.cancer.gov/csr/1975_2008/.
11. Zebrack B, Mathews-Bradshaw B, Siegel S. Quality cancer care for adolescents and young adults: a position statement. *J Clin Oncol* 2010;28:4862-4867. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20855821>.
12. Alman BA, Pajerski ME, Diaz-Cano S, et al. Aggressive fibromatosis (desmoid tumor) is a monoclonal disorder. *Diagn Mol Pathol* 1997;6:98-101. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9098648>.
13. Abramson DH, Ellsworth RM, Kitchin FD, Tung G. Second nonocular tumors in retinoblastoma survivors. Are they radiation-induced? *Ophthalmology* 1984;91:1351-1355. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6595610>.
14. Malkin D, Li FP, Strong LC, et al. Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science* 1990;250:1233-1238. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1978757>.
15. Miettinen M, Wang ZF, Sarlomo-Rikala M, et al. Succinate dehydrogenase-deficient GISTs: a clinicopathologic, immunohistochemical, and molecular genetic study of 66 gastric GISTs with predilection to young age. *Am J Surg Pathol* 2011;35:1712-1721. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21997692>.



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

16. Janeway KA, Kim SY, Lodish M, et al. Defects in succinate dehydrogenase in gastrointestinal stromal tumors lacking KIT and PDGFRA mutations. *Proc Natl Acad Sci U S A* 2011;108:314-318. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21173220>.

17. Friedman JM. Neurofibromatosis 1. In: Pagon RA, Bird TD, Dolan CR, Stephens K, eds. *GeneReviews* [Internet]. Seattle, WA: University of Washington, Seattle, WA. Initial Posting: October 2, 1998; Last Update: June 2, 2009. <http://www.ncbi.nlm.nih.gov/books/NBK1109/>.

18. Smith RA, Cokkinides V, Brooks D, et al. Cancer screening in the United States, 2010: a review of current American Cancer Society guidelines and issues in cancer screening. *CA Cancer J Clin* 2010;60:99-9119. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20228384>.

19. Bryant H. Screening for cancer in children, adolescents, and young adults: questions-and more questions. *Cancer* 2011;117:2275-2280. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523746>.

20. Pollock BH, Krischer JP, Vietti TJ. Interval between symptom onset and diagnosis of pediatric solid tumors. *J Pediatr* 1991;119:725-732. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1941378>.

21. Klein-Geltink J, Pogany L, Mery LS, et al. Impact of age and diagnosis on waiting times between important healthcare events among children 0 to 19 years cared for in pediatric units: the Canadian Childhood Cancer Surveillance and Control Program. *J Pediatr Hematol Oncol* 2006;28:433-439. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16825989>.

22. Dang-Tan T, Trottier H, Mery LS, et al. Delays in diagnosis and treatment among children and adolescents with cancer in Canada. *Pediatr Blood Cancer* 2008;51:468-474. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18454472>.

23. Martin S, Ulrich C, Munsell M, et al. Delays in cancer diagnosis in underinsured young adults and older adolescents. *Oncologist*

2007;12:816-824. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17673613>.

24. Ferrari A, Thomas D, Franklin AR, et al. Starting an adolescent and young adult program: some success stories and some obstacles to overcome. *J Clin Oncol* 2010;28:4850-4857. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20479411>.

25. Boissel N, Auclerc M-F, Lheritier V, et al. Should adolescents with acute lymphoblastic leukemia be treated as old children or young adults? Comparison of the French FRALLE-93 and LALA-94 trials. *J Clin Oncol* 2003;21:774-780. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12610173>.

26. de Bont JM, Holt Bvd, Dekker AW, et al. Significant difference in outcome for adolescents with acute lymphoblastic leukemia treated on pediatric vs adult protocols in the Netherlands. *Leukemia* 2004;18:2032-2035. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15483674>.

27. Ribera J-M, Oriol A, Sanz M-A, et al. Comparison of the results of the treatment of adolescents and young adults with standard-risk acute lymphoblastic leukemia with the Programa Espanol de Tratamiento en Hematologia pediatric-based protocol ALL-96. *J Clin Oncol* 2008;26:1843-1849. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18398150>.

28. Stock W, La M, Sanford B, et al. What determines the outcomes for adolescents and young adults with acute lymphoblastic leukemia treated on cooperative group protocols? A comparison of Children's Cancer Group and Cancer and Leukemia Group B studies. *Blood* 2008;112:1646-1654. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18502832>.

29. Huguet F, Leguay T, Raffoux E, et al. Pediatric-inspired therapy in adults with Philadelphia chromosome-negative acute lymphoblastic leukemia: the GRAALL-2003 study. *J Clin Oncol* 2009;27:911-918. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19124805>.

30. Ferrari A, Dileo P, Casanova M, et al. Rhabdomyosarcoma in adults. A retrospective analysis of 171 patients treated at a single institution. *Cancer* 2003;98:571-580. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12879475>.

31. Scurr M, Judson I. How to treat the Ewing's family of sarcomas in adult patients. *Oncologist* 2006;11:65-72. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16401715>.

32. Sandlund JT. Should adolescents with NHL be treated as old children or young adults? *Hematology Am Soc Hematol Educ Program* 2007:297-303. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18024643>.

33. Tai E, Pollack LA, Townsend J, et al. Differences in non-Hodgkin lymphoma survival between young adults and children. *Arch Pediatr Adolesc Med* 2010;164:218-224. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20194253>.

34. Burkhardt B, Oschlies I, Klapper W, et al. Non-Hodgkin's lymphoma in adolescents: experiences in 378 adolescent NHL patients treated according to pediatric NHL-BFM protocols. *Leukemia* 2011;25:153-160. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21030984>.

35. Bleyer WA, Tejada H, Murphy SB, et al. National cancer clinical trials: children have equal access; adolescents do not. *J Adolesc Health* 1997;21:366-373. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9401854>.

36. Shochat SJ, Fremgen AM, Murphy SB, et al. Childhood cancer: patterns of protocol participation in a national survey. *CA Cancer J Clin* 2001;51:119-130. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11577480>.

37. Downs-Canner S, Shaw PH. A comparison of clinical trial enrollment between adolescent and young adult (AYA) oncology patients treated at affiliated adult and pediatric oncology centers. *J*

Pediatr Hematol Oncol 2009;31:927-929. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19855302>.

38. Parsons HM, Harlan LC, Seibel NL, et al. Clinical trial participation and time to treatment among adolescents and young adults with cancer: does age at diagnosis or insurance make a difference? *J Clin Oncol* 2011;29:4045-4053. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21931022>.

39. Howell DL, Ward KC, Austin HD, et al. Access to pediatric cancer care by age, race, and diagnosis, and outcomes of cancer treatment in pediatric and adolescent patients in the state of Georgia. *J Clin Oncol* 2007;25:4610-4615. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17925556>.

40. Albritton KH, Wiggins CH, Nelson HE, Weeks JC. Site of oncologic specialty care for older adolescents in Utah. *J Clin Oncol* 2007;25:4616-4621. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17925557>.

41. Bleyer A. The Quid Pro Quo of pediatric versus adult services for older adolescent cancer patients. *Pediatr Blood Cancer* 2010;54:238-241. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19813248>.

42. Veal GJ, Hartford CM, Stewart CF. Clinical pharmacology in the adolescent oncology patient. *J Clin Oncol* 2010;28:4790-4799. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20439647>.

43. Oeffinger KC, Mertens AC, Hudson MM, et al. Health care of young adult survivors of childhood cancer: a report from the Childhood Cancer Survivor Study. *Ann Fam Med* 2004;2:61-70. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15053285>.

44. Oeffinger KC, Hudson MM. Long-term complications following childhood and adolescent cancer: foundations for providing risk-based health care for survivors. *CA Cancer J Clin* 2004;54:208-236. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15253918>.



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

45. Armstrong GT, Stovall M, Robison LL. Long-term effects of radiation exposure among adult survivors of childhood cancer: results from the childhood cancer survivor study. *Radiat Res* 2010;174:840-850. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21128808>.
46. Levine J, Canada A, Stern CJ. Fertility preservation in adolescents and young adults with cancer. *J Clin Oncol* 2010;28:4831-4841. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20458029>.
47. Janson C, Leisenring W, Cox C, et al. Predictors of marriage and divorce in adult survivors of childhood cancers: a report from the Childhood Cancer Survivor Study. *Cancer Epidemiol Biomarkers Prev* 2009;18:2626-2635. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19815636>.
48. Bleyer A, Choi M, Wang SJ, et al. Increased vulnerability of the spinal cord to radiation or intrathecal chemotherapy during adolescence: A report from the Children's Oncology Group. *Pediatr Blood Cancer* 2009;53:1205-1210. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19821538>.
49. Hijjiya N, Ness KK, Ribeiro RC, Hudson MM. Acute leukemia as a secondary malignancy in children and adolescents: current findings and issues. *Cancer* 2009;115:23-35. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19072983>.
50. Knight KR, Kraemer DF, Neuwelt EA. Ototoxicity in children receiving platinum chemotherapy: underestimating a commonly occurring toxicity that may influence academic and social development. *J Clin Oncol* 2005;23:8588-8596. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16314621>.
51. Schell MJ, McHaney VA, Green AA, et al. Hearing loss in children and young adults receiving cisplatin with or without prior cranial irradiation. *J Clin Oncol* 1989;7:754-760. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2715805>.
52. Skinner R, Cotterill SJ, Stevens MC. Risk factors for nephrotoxicity after ifosfamide treatment in children: a UKCCSG Late Effects Group study. United Kingdom Children's Cancer Study Group. *Br J Cancer* 2000;82:1636-1645. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10817497>.
53. Roscoe JA, Morrow GR, Apro MS, et al. Anticipatory nausea and vomiting. *Support Care Cancer* 2011;19:1533-1538. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20803345>.
54. Figueroa-Moseley C, Jean-Pierre P, Roscoe JA, et al. Behavioral interventions in treating anticipatory nausea and vomiting. *J Natl Compr Canc Netw* 2007;5:44-50. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17239325>.
55. Armenian SH, Sun CL, Kawashima T, et al. Long-term health-related outcomes in survivors of childhood cancer treated with hematopoietic cell transplantation (HCT) versus conventional therapy: a report from the Bone Marrow Transplant Survivor Study (BMTSS) and Childhood Cancer Survivor Study (CCSS). *Blood* 2011. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21652685>.
56. Rizzo JD, Curtis RE, Socie G, et al. Solid cancers after allogeneic hematopoietic cell transplantation. *Blood* 2009;113:1175-1183. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18971419>.
57. Wallace WH. Oncofertility and preservation of reproductive capacity in children and young adults. *Cancer* 2011;117:2301-2310. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523750>.
58. Lee SJ, Schover LR, Partridge AH, et al. American Society of Clinical Oncology recommendations on fertility preservation in cancer patients. *J Clin Oncol* 2006;24:2917-2931. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16651642>.
59. Wallace WH, Anderson RA, Irvine DS. Fertility preservation for young patients with cancer: who is at risk and what can be offered?



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

Lancet Oncol 2005;6:209-218. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15811616>.

60. Darzy KH. Radiation-induced hypopituitarism after cancer therapy: who, how and when to test. Nat Clin Pract Endocrinol Metab 2009;5:88-99. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19165221>.

61. Redig AJ, Brannigan R, Stryker SJ, et al. Incorporating fertility preservation into the care of young oncology patients. Cancer 2011;117:4-10. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21235031>.

62. Green DM, Kawashima T, Stovall M, et al. Fertility of female survivors of childhood cancer: a report from the childhood cancer survivor study. J Clin Oncol 2009;27:2677-2685. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19364965>.

63. Green DM, Nolan VG, Kawashima T, et al. Decreased fertility among female childhood cancer survivors who received 22-27 Gy hypothalamic/pituitary irradiation: a report from the Childhood Cancer Survivor Study. Fertil Steril 2011;95:1922-1927, 1927 e1921. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21376314>.

64. Chemaitilly W, Mertens AC, Mitby P, et al. Acute ovarian failure in the childhood cancer survivor study. J Clin Endocrinol Metab 2006;91:1723-1728. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16492690>.

65. Green DM, Sklar CA, Boice JD, Jr., et al. Ovarian failure and reproductive outcomes after childhood cancer treatment: results from the Childhood Cancer Survivor Study. J Clin Oncol 2009;27:2374-2381. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19364956>.

66. Wo JY, Viswanathan AN. Impact of radiotherapy on fertility, pregnancy, and neonatal outcomes in female cancer patients. Int J Radiat Oncol Biol Phys 2009;73:1304-1312. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19306747>.

67. Lobo RA. Potential options for preservation of fertility in women. N Engl J Med 2005;353:64-73. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16000356>.

68. Bines J, Oleske DM, Cobleigh MA. Ovarian function in premenopausal women treated with adjuvant chemotherapy for breast cancer. J Clin Oncol 1996;14:1718-1729. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/8622093>.

69. Goodwin PJ, Ennis M, Pritchard KI, et al. Risk of menopause during the first year after breast cancer diagnosis. J Clin Oncol 1999;17:2365-2370. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10561298>.

70. De Bruin ML, Huisbrink J, Hauptmann M, et al. Treatment-related risk factors for premature menopause following Hodgkin lymphoma. Blood 2008;111:101-108. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17890454>.

71. Haukvik UKH, Dieset I, Bjoro T, et al. Treatment-related premature ovarian failure as a long-term complication after Hodgkin's lymphoma. Ann Oncol 2006;17:1428-1433. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16831852>.

72. Terenziani M, Piva L, Meazza C, et al. Oophoropexy: a relevant role in preservation of ovarian function after pelvic irradiation. Fertil Steril 2009;91:935 e915-936. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18951125>.

73. Badawy A, Elnashar A, El-Ashry M, Shahat M. Gonadotropin-releasing hormone agonists for prevention of chemotherapy-induced ovarian damage: prospective randomized study. Fertil Steril 2009;91:694-697. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18675959>.

74. Behringer K, Wildt L, Mueller H, et al. No protection of the ovarian follicle pool with the use of GnRH-analogues or oral contraceptives in young women treated with escalated BEACOPP for advanced-stage Hodgkin lymphoma. Final results of a phase II trial from the German



Hodgkin Study Group. *Ann Oncol* 2010;21:2052-2060. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20305034>.

75. Bedaiwy MA, Abou-Setta AM, Desai N, et al. Gonadotropin-releasing hormone analog cotreatment for preservation of ovarian function during gonadotoxic chemotherapy: a systematic review and meta-analysis. *Fertil Steril* 2011;95:906-914 e901-904. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21145541>.

76. Azim H, Jr., Gentilini O, Locatelli M, et al. Managing pregnant women with cancer: personal considerations and a review of the literature. *Ecancermedicalscience* 2011;5:204. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22276051>.

77. Azim HA, Jr., Pavlidis N, Peccatori FA. Treatment of the pregnant mother with cancer: a systematic review on the use of cytotoxic, endocrine, targeted agents and immunotherapy during pregnancy. Part II: Hematological tumors. *Cancer Treat Rev* 2010;36:110-121. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20018452>.

78. Azim HA, Jr., Peccatori FA, Pavlidis N. Treatment of the pregnant mother with cancer: a systematic review on the use of cytotoxic, endocrine, targeted agents and immunotherapy during pregnancy. Part I: Solid tumors. *Cancer Treat Rev* 2010;36:101-109. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20015593>.

79. Chemaitilly W, Sklar CA. Endocrine complications in long-term survivors of childhood cancers. *Endocr Relat Cancer* 2010;17:R141-159. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20453080>.

80. Kenney LB, Laufer MR, Grant FD, et al. High risk of infertility and long term gonadal damage in males treated with high dose cyclophosphamide for sarcoma during childhood. *Cancer* 2001;91:613-621. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11169946>.

81. Lee SJ, Schover LR, Partridge AH, et al. American Society of Clinical Oncology recommendations on fertility preservation in cancer

patients. *J Clin Oncol* 2006;24:2917-2931. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16651642>.

82. Zebrack BJ. Psychological, social, and behavioral issues for young adults with cancer. *Cancer* 2011;117:2289-2294. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523748>.

83. Abrams AN, Hazen EP, Penson RT. Psychosocial issues in adolescents with cancer. *Cancer Treat Rev* 2007;33:622-630. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17434265>.

84. Zebrack B, Isaacson S. Psychosocial Care of Adolescent and Young Adult (AYA) Cancer Patients and Survivors. *J Clin Oncol* 2011:In press. Available at:

85. D'Agostino NM, Penney A, Zebrack B. Providing developmentally appropriate psychosocial care to adolescent and young adult cancer survivors. *Cancer* 2011;117:2329-2334. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523754>.

86. Zeltzer LK. Cancer in adolescents and young adults psychosocial aspects. Long-term survivors. *Cancer* 1993;71:3463-3468. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8490896>.

87. Arnett JJ. Emerging adulthood. A theory of development from the late teens through the twenties. *Am Psychol* 2000;55:469-480. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10842426>.

88. Morgan S, Davies S, Palmer S, Plaster M. Sex, drugs, and rock 'n' roll: caring for adolescents and young adults with cancer. *J Clin Oncol* 2010;28:4825-4830. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20498401>.

89. Vinokur AD, Threatt BA, Vinokur-Kaplan D, Satariano WA. The process of recovery from breast cancer for younger and older patients. Changes during the first year. *Cancer* 1990;65:1242-1254. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2302673>.



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

90. Mor V, Allen S, Malin M. The psychosocial impact of cancer on older versus younger patients and their families. *Cancer* 1994;74:2118-2127. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8087779>.
91. Stava CJ, Lopez A, Vassilopoulou-Sellin R. Health profiles of younger and older breast cancer survivors. *Cancer* 2006;107:1752-1759. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16967441>.
92. Kazak AE, Derosa BW, Schwartz LA, et al. Psychological outcomes and health beliefs in adolescent and young adult survivors of childhood cancer and controls. *J Clin Oncol* 2010;28:2002-2007. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20231679>.
93. Krull KR, Huang S, Gurney JG, et al. Adolescent behavior and adult health status in childhood cancer survivors. *J Cancer Surviv* 2010;4:210-217. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20383785>.
94. Emmons K, Li FP, Whitton J, et al. Predictors of smoking initiation and cessation among childhood cancer survivors: a report from the childhood cancer survivor study. *J Clin Oncol* 2002;20:1608-1616. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11896111>.
95. Lown EA, Goldsby R, Mertens AC, et al. Alcohol consumption patterns and risk factors among childhood cancer survivors compared to siblings and general population peers. *Addiction* 2008;103:1139-1148. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18554347>.
96. Cox CL, McLaughlin RA, Steen BD, Hudson MM. Predicting and modifying substance use in childhood cancer survivors: application of a conceptual model. *Oncol Nurs Forum* 2006;33:51-60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16470234>.
97. Zebrack B, Bleyer A, Albritton K, et al. Assessing the health care needs of adolescent and young adult cancer patients and survivors. *Cancer* 2006;107:2915-2923. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17103383>.
98. Palefsky JM, Giuliano AR, Goldstone S, et al. HPV vaccine against anal HPV infection and anal intraepithelial neoplasia. *N Engl J Med* 2011;365:1576-1585. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22029979>.
99. Butow P, Palmer S, Pai A, et al. Review of adherence-related issues in adolescents and young adults with cancer. *J Clin Oncol* 2010;28:4800-4809. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20212260>.
100. Kondryn HJ, Edmondson CL, Hill J, Eden TO. Treatment non-adherence in teenage and young adult patients with cancer. *Lancet Oncol* 2011;12:100-108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20580606>.
101. Windebank KP, Spinetta JJ. Do as I say or die: compliance in adolescents with cancer. *Pediatr Blood Cancer* 2008;50:1099-1100. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18360837>.
102. Kato PM, Cole SW, Bradlyn AS, Pollock BH. A video game improves behavioral outcomes in adolescents and young adults with cancer: a randomized trial. *Pediatrics* 2008;122:e305-317. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18676516>.
103. Kahana S, Drotar D, Frazier T. Meta-analysis of psychological interventions to promote adherence to treatment in pediatric chronic health conditions. *J Pediatr Psychol* 2008;33:590-611. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18192300>.
104. Albritton K, Barr R, Bleyer A. The adolescence of young adult oncology. *Semin Oncol* 2009;36:478-488. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19835743>.
105. Kyngas H, Mikkonen R, Nousiainen EM, et al. Coping with the onset of cancer: coping strategies and resources of young people with cancer. *Eur J Cancer Care (Engl)* 2001;10:6-11. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11827269>.

106. Zebrack BJ, Oeffinger KC, Hou P, Kaplan S. Advocacy skills training for young adult cancer survivors: the Young Adult Survivors Conference at Camp Mak-a-Dream. Support Care Cancer 2006;14:779-782. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16482447>.

107. Treadgold CL, Kuperberg A. Been there, done that, wrote the blog: the choices and challenges of supporting adolescents and young adults with cancer. J Clin Oncol 2010;28:4842-4849. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20351337>.

108. Zebrack B. Information and service needs for young adult cancer patients. Support Care Cancer 2008;16:1353-1360. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18386075>.

109. Zebrack B. Information and service needs for young adult cancer survivors. Support Care Cancer 2009;17:349-357. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18543006>.

110. Palmer S, Mitchell A, Thompson K, Sexton M. Unmet needs among adolescent cancer patients: a pilot study. Palliat Support Care 2007;5:127-134. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17578063>.

111. Fallon S, Smith J, Morgan S, et al. 'Pizza, patients and points of view': Involving young people in the design of a post registration module entitled the adolescent with cancer. Nurse Educ Pract 2008;8:140-147. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17919977>.

112. Rabin C, Simpson N, Morrow K, Pinto B. Behavioral and psychosocial program needs of young adult cancer survivors. Qual Health Res 2011;21:796-806. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20705863>.

113. Kirchhoff AC, Leisenring W, Krull KR, et al. Unemployment among adult survivors of childhood cancer: a report from the childhood cancer

survivor study. Med Care 2010;48:1015-1025. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20940653>.

114. Zeltzer LK, Recklitis C, Buchbinder D, et al. Psychological status in childhood cancer survivors: a report from the Childhood Cancer Survivor Study. J Clin Oncol 2009;27:2396-2404. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19255309>.

115. Soliman H, Agresta SV. Current issues in adolescent and young adult cancer survivorship. Cancer Control 2008;15:55-62. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18094661>.

116. Oeffinger KC, Tonorezos ES. The cancer is over, now what?: Understanding risk, changing outcomes. Cancer 2011;117:2250-2257. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523742>.

117. Woodward E, Jessop M, Glaser A, Stark D. Late effects in survivors of teenage and young adult cancer: does age matter? Ann Oncol 2011;22 2561-2568. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21427066>.

118. Oeffinger KC, Mertens AC, Sklar CA, et al. Chronic health conditions in adult survivors of childhood cancer. N Engl J Med 2006;355:1572-1582. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17035650>.

119. Armstrong GT, Liu Q, Yasui Y, et al. Late mortality among 5-year survivors of childhood cancer: a summary from the Childhood Cancer Survivor Study. J Clin Oncol 2009;27:2328-2338. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19332714>.

120. Dores GM, Metayer C, Curtis RE, et al. Second malignant neoplasms among long-term survivors of Hodgkin's disease: a population-based evaluation over 25 years. J Clin Oncol 2002;20:3484-3494. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12177110>.

121. Aleman BM, van den Belt-Dusebout AW, De Bruin ML, et al. Late cardiotoxicity after treatment for Hodgkin lymphoma. Blood



2007;109:1878-1886. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17119114>.

122. Curtis RE, Freedman DM, Ron E, et al. New Malignancies Among Cancer Survivors: SEER Cancer Registries, 1973-2000. National Cancer Institute, NIH Publ. No. 05-5302. Bethesda, MD, 2006. 2006. Available at:

http://seer.cancer.gov/publications/mpmono/MPMonograph_complete.pdf.

123. Ng AK, Travis LB. Subsequent malignant neoplasms in cancer survivors. *Cancer J* 2008;14:429-434. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19060610>.

124. Ng AK, Kenney LB, Gilbert ES, Travis LB. Secondary malignancies across the age spectrum. *Semin Radiat Oncol* 2010;20:67-78. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19959033>.

125. Friedman DL, Whitton J, Leisenring W, et al. Subsequent neoplasms in 5-year survivors of childhood cancer: the Childhood Cancer Survivor Study. *J Natl Cancer Inst* 2010;102:1083-1095. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20634481>.

126. Meadows AT, Friedman DL, Neglia JP, et al. Second neoplasms in survivors of childhood cancer: findings from the Childhood Cancer Survivor Study cohort. *J Clin Oncol* 2009;27:2356-2362. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19255307>.

127. Travis LB, Hill DA, Dores GM, et al. Breast cancer following radiotherapy and chemotherapy among young women with Hodgkin disease. *JAMA* 2003;290:465-475. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12876089>.

128. van Leeuwen FE, Klokman WJ, Stovall M, et al. Roles of radiation dose, chemotherapy, and hormonal factors in breast cancer following Hodgkin's disease. *J Natl Cancer Inst* 2003;95:971-980. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12837833>.

129. Travis LB, Hill D, Dores GM, et al. Cumulative absolute breast cancer risk for young women treated for Hodgkin lymphoma. *J Natl Cancer Inst* 2005;97:1428-1437. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16204692>.

130. Henderson TO, Amsterdam A, Bhatia S, et al. Systematic review: surveillance for breast cancer in women treated with chest radiation for childhood, adolescent, or young adult cancer. *Ann Intern Med* 2010;152:444-455; W144-454. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20368650>.

131. Travis LB, Gospodarowicz M, Curtis RE, et al. Lung cancer following chemotherapy and radiotherapy for Hodgkin's disease. *J Natl Cancer Inst* 2002;94:182-192. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11830608>.

132. Swerdlow AJ, Higgins CD, Smith P, et al. Second Cancer Risk After Chemotherapy for Hodgkin's Lymphoma: A Collaborative British Cohort Study. *J Clin Oncol* 2011;29:4096-4104. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21969511>.

133. Travis LB, Beard C, Allan JM, et al. Testicular cancer survivorship: research strategies and recommendations. *J Natl Cancer Inst* 2010;102:1114-1130. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20585105>.

134. Gilligan T. Testicular cancer survivorship. *Hematol Oncol Clin North Am* 2011;25:627-639, x. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21570614>.

135. Fossa SD, Chen J, Schonfeld SJ, et al. Risk of contralateral testicular cancer: a population-based study of 29,515 U.S. men. *J Natl Cancer Inst* 2005;97:1056-1066. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16030303>.

136. Travis LB, Fossa SD, Schonfeld SJ, et al. Second cancers among 40,576 testicular cancer patients: focus on long-term survivors. *J Natl*



Cancer Inst 2005;97:1354-1365. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16174857>.

137. Wierecky J, Kollmannsberger C, Boehlke I, et al. Secondary leukemia after first-line high-dose chemotherapy for patients with advanced germ cell cancer. *J Cancer Res Clin Oncol* 2005;131:255-260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15627215>.

138. Howard R, Gilbert E, Lynch CF, et al. Risk of leukemia among survivors of testicular cancer: a population-based study of 42,722 patients. *Ann Epidemiol* 2008;18:416-421. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18433667>.

139. Chaturvedi AK, Engels EA, Gilbert ES, et al. Second cancers among 104,760 survivors of cervical cancer: evaluation of long-term risk. *J Natl Cancer Inst* 2007;99:1634-1643. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17971527>.

140. Brown LM, Chen BE, Pfeiffer RM, et al. Risk of second non-hematological malignancies among 376,825 breast cancer survivors. *Breast Cancer Res Treat* 2007;106:439-451. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17277968>.

141. Hemminki K, Lenner P, Sundquist J, Bermejo JL. Risk of subsequent solid tumors after non-Hodgkin's lymphoma: effect of diagnostic age and time since diagnosis. *J Clin Oncol* 2008;26:1850-1857. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18347006>.

142. Ginsberg JP, Goodman P, Leisenring W, et al. Long-term survivors of childhood Ewing sarcoma: report from the childhood cancer survivor study. *J Natl Cancer Inst* 2010;102:1272-1283. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20656964>.

143. Nagarajan R, Kamruzzaman A, Ness KK, et al. Twenty years of follow-up of survivors of childhood osteosarcoma: a report from the Childhood Cancer Survivor Study. *Cancer* 2011;117:625-634. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20922787>.

144. Children's Oncology Group. Long-term follow-up guidelines for survivors of childhood, adolescent and young adult cancers, Version 3.0.; 2008. Available at: www.survivorshipguidelines.org.

145. Chen MH, Colan SD, Diller L. Cardiovascular disease: cause of morbidity and mortality in adult survivors of childhood cancers. *Circ Res* 2011;108:619-628. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21372293>.

146. Reinders JG, Heijmen BJ, Olofsen-van Acht MJ, et al. Ischemic heart disease after mantlefield irradiation for Hodgkin's disease in long-term follow-up. *Radiother Oncol* 1999;51:35-42. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10386715>.

147. Swerdlow AJ, Higgins CD, Smith P, et al. Myocardial infarction mortality risk after treatment for Hodgkin disease: a collaborative British cohort study. *J Natl Cancer Inst* 2007;99:206-214. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17284715>.

148. van den Belt-Dusebout AW, Nuver J, de Wit R, et al. Long-term risk of cardiovascular disease in 5-year survivors of testicular cancer. *J Clin Oncol* 2006;24:467-475. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16421423>.

149. Haugnes HS, Wethal T, Aass N, et al. Cardiovascular risk factors and morbidity in long-term survivors of testicular cancer: a 20-year follow-up study. *J Clin Oncol* 2010;28:4649-4657. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20855830>.

150. Abouassaly R, Fossa SD, Giwercman A, et al. Sequelae of treatment in long-term survivors of testis cancer. *Eur Urol* 2011;60:516-526. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21684072>.

151. Mulrooney DA, Yeazel MW, Kawashima T, et al. Cardiac outcomes in a cohort of adult survivors of childhood and adolescent cancer: retrospective analysis of the Childhood Cancer Survivor Study cohort. *BMJ* 2009;339:b4606. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19996459>.



NCCN Guidelines Version 1.2012

Adolescent and Young Adult Oncology

152. Mertens AC, Yasui Y, Liu Y, et al. Pulmonary complications in survivors of childhood and adolescent cancer. A report from the Childhood Cancer Survivor Study. *Cancer* 2002;95:2431-2441. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12436452>.

153. Fossa SD, Gilbert E, Dores GM, et al. Noncancer causes of death in survivors of testicular cancer. *J Natl Cancer Inst* 2007;99:533-544. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17405998>.

154. O'Sullivan JM, Huddart RA, Norman AR, et al. Predicting the risk of bleomycin lung toxicity in patients with germ-cell tumours. *Ann Oncol* 2003;14:91-96. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12488299>.

155. Hagnes HS, Aass N, Fossa SD, et al. Pulmonary function in long-term survivors of testicular cancer. *J Clin Oncol* 2009;27:2779-2786. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19414680>.

156. Diller L, Chow EJ, Gurney JG, et al. Chronic disease in the Childhood Cancer Survivor Study cohort: a review of published findings. *J Clin Oncol* 2009;27:2339-2355. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19364955>.

157. Packer RJ, Gurney JG, Punyko JA, et al. Long-term neurologic and neurosensory sequelae in adult survivors of a childhood brain tumor: childhood cancer survivor study. *J Clin Oncol* 2003;21:3255-3261. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12947060>.

158. Brydoy M, Oldenburg J, Klepp O, et al. Observational study of prevalence of long-term Raynaud-like phenomena and neurological side effects in testicular cancer survivors. *J Natl Cancer Inst* 2009;101:1682-1695. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19940282>.

159. Bowers DC, McNeil DE, Liu Y, et al. Stroke as a late treatment effect of Hodgkin's Disease: a report from the Childhood Cancer Survivor Study. *J Clin Oncol* 2005;23:6508-6515. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16170160>.

160. Bowers DC, Liu Y, Leisenring W, et al. Late-occurring stroke among long-term survivors of childhood leukemia and brain tumors: a report from the Childhood Cancer Survivor Study. *J Clin Oncol* 2006;24:5277-5282. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17088567>.

161. De Bruin ML, Dorresteijn LD, van't Veer MB, et al. Increased risk of stroke and transient ischemic attack in 5-year survivors of Hodgkin lymphoma. *J Natl Cancer Inst* 2009;101:928-937. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19535773>.

162. Chow EJ, Friedman DL, Stovall M, et al. Risk of thyroid dysfunction and subsequent thyroid cancer among survivors of acute lymphoblastic leukemia: a report from the Childhood Cancer Survivor Study. *Pediatr Blood Cancer* 2009;53:432-437. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19459201>.

163. Bhatti P, Veiga LH, Ronckers CM, et al. Risk of second primary thyroid cancer after radiotherapy for a childhood cancer in a large cohort study: an update from the childhood cancer survivor study. *Radiat Res* 2010;174:741-752. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21128798>.

164. Huddart RA, Norman A, Moynihan C, et al. Fertility, gonadal and sexual function in survivors of testicular cancer. *Br J Cancer* 2005;93:200-207. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15999104>.

165. Shankar SM, Marina N, Hudson MM, et al. Monitoring for cardiovascular disease in survivors of childhood cancer: report from the Cardiovascular Disease Task Force of the Children's Oncology Group. *Pediatrics* 2008;121:e387-396. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18187811>.

166. Liles A, Blatt J, Morris D, et al. Monitoring pulmonary complications in long-term childhood cancer survivors: guidelines for the primary care physician. *Cleve Clin J Med* 2008;75:531-539. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18646589>.



167. Freyer DR. Transition of care for young adult survivors of childhood and adolescent cancer: rationale and approaches. *J Clin Oncol* 2010;28:4810-4818. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20351333>.

168. Nathan PC, Hayes-Lattin B, Sisler JJ, Hudson MM. Critical issues in transition and survivorship for adolescents and young adults with cancers. *Cancer* 2011;117:2335-2341. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523755>.

169. Oeffinger KC, McCabe MS. Models for delivering survivorship care. *J Clin Oncol* 2006;24:5117-5124. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17093273>.

170. Snyder CF, Earle CC, Herbert RJ, et al. Preventive care for colorectal cancer survivors: a 5-year longitudinal study. *J Clin Oncol* 2008;26:1073-1079. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18309941>.

171. Snyder CF, Frick KD, Kantsiper ME, et al. Prevention, screening, and surveillance care for breast cancer survivors compared with controls: changes from 1998 to 2002. *J Clin Oncol* 2009;27:1054-1061. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19164212>.

172. Blaauwbroek R, Tuinier W, Meyboom-de Jong B, et al. Shared care by paediatric oncologists and family doctors for long-term follow-up of adult childhood cancer survivors: a pilot study. *Lancet Oncol* 2008;9:232-238. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18282804>.

173. Wallace WH, Blacklay A, Eiser C, et al. Developing strategies for long term follow up of survivors of childhood cancer. *BMJ* 2001;323:271-274. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11485960>.

174. Oeffinger KC. Longitudinal risk-based health care for adult survivors of childhood cancer. *Curr Probl Cancer* 2003;27:143-167. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12748583>.

175. Wein S, Pery S, Zer A. Role of palliative care in adolescent and young adult oncology. *J Clin Oncol* 2010;28:4819-4824. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20212259>.

176. Pritchard S, Cuvelier G, Harlos M, Barr R. Palliative care in adolescents and young adults with cancer. *Cancer* 2011;117:2323-2328. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523753>.

177. Lyon ME, McCabe MA, Patel KM, D'Angelo LJ. What do adolescents want? An exploratory study regarding end-of-life decision-making. *J Adolesc Health* 2004;35:529 e521-526. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15581537>.

178. Webb NM, Tucker D. Young adults' opinions about hospice and home death. *J Palliat Med* 2009;12:337-342. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19327070>.

179. Bell CJ, Skiles J, Pradhan K, Champion VL. End-of-life experiences in adolescents dying with cancer. *Support Care Cancer* 2010;18:827-835. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19727847>.

180. Wiener L, Ballard E, Brennan T, et al. How I wish to be remembered: the use of an advance care planning document in adolescent and young adult populations. *J Palliat Med* 2008;11:1309-1313. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19115889>.