ACTEMRA® Infusion Referral Form



4510 Richmond Road Warrensville Heights, OH 44128

Phone: 800-552-8442 Fax: 216-201-5127

Please complete each section of the referral form below and fax along with a copy (front and back) of <u>all</u> of the patient's pharmacy and medical insurance cards, the patient's demographic face sheet, and any relevant clinical notes/documents.							
Prescriber Info.	Prescriber:				NPI:		
	Phone: Office Contact: Office Contact:						
Patient Information	Name:				DOB: □M □F		
	Address:						
	Phone:		2 nd Phone:		MRN:		
	Primary Language: Functional Limitation				·		
Clinical Information	Diagnosis (Include ICD-10 Code):						
	Weight: □ lb □ kg Height: □ in				IV access: □PIV □PICC □Port □Other:		
	Patient's first dose of IV Actemra? Yes No (If no, date of last dose: Prior dose (in mg):)						
	Allergies:Latex allergy? Yes No						
	Prior treatments/Reason for discontinuation:						
	Date of <u>negative</u> TB test: TB test pending, will fax results. Patient is HBV negative or has been treated: _Yes _No						
	History of kidney disease: ☐Yes ☐No If yes, SCr: GFR/CrCl: History of heart failure: ☐Yes ☐No						
	Required Labs:	ANC	Platelets AST			ALT	SCr
	Result:		Result: Upper limit of normal			Result:	
	nesuit.				l:	Upper limit of normal:	
	Date:						
	Referring provider's preferred site of care: Home Care Infusion Center Home Infusion Home Care to determine site of care						
	Site of care preference is subject to payer limitations, clinical appropriateness, and the availability of servicing providers. Based on the clinical judgement of the pharmacist, doses may be rounded up or down by no more than 10% unless checked here:						
Prescription Information	Actemra® Dose		Infusion Diluent/Volume		Rate	Frequency	Number of Doses
	Adult Rheumatoid Arthritis*		in 100mL NaCl 0.9%		Infused over		
	□4mg/kg				60 minutes	every four weeks	
	□8mg/kg Polyarticular JIA		Weight <30kg: in 50mL NaCl 0.9% Weight ≥30kg: in 100mL NaCl 0.9%		Infused over		
	☐ 10mg/kg (weight <30kg)				60 minutes	every four weeks	
	☐ 8mg/kg (weight ≥30kg) Systemic JIA						
	□ 12mg/kg (weight <30kg)		Weight <30kg: in 50mL NaCl 0.9% Weight ≥30kg: in 100mL NaCl 0.9%		Infused over	every two weeks	
	□8mg/kg (weight ≥30kg)				60 minutes		
	*Doses exceeding 800mg per infusion are not recommended.						
	Premedication/PRN medication orders:						
	Laboratory orders: □ANC/Platelets/AST/ALT 4 to 8 weeks after the start of therapy and every 3 months thereafter.						
	Other lab orders (subject to availability):						
Prescriber Signature	My signature for this prescription also confirms that the treatment(s) indicated on this referral is/are medically necessary. I authorize Home						
	Care and its representatives to act as an agent of mine to initiate and execute the patient's insurance prior authorization process and to provide infusion-related nursing services and supplies in conjunction with the therapy prescribed above.						
	Circustums.						
	Signature:					Date:	